

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

Jean-Charles Preiser
Editor

 Springer

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

Jean-Charles Preiser

Editor

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

 Springer

Editor
Jean-Charles Preiser
Brussels
Belgium

ISBN 978-3-319-27685-4 ISBN 978-3-319-27687-8 (eBook)
DOI 10.1007/978-3-319-27687-8

Library of Congress Control Number: 2016935706

Springer Cham Heidelberg New York Dordrecht London
© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer International Publishing AG Switzerland is part of Springer Science+Business Media
(www.springer.com)

Foreword

Life is like riding a bicycle. To keep your balance, you must keep moving.

Albert Einstein

All living organisms share common biological characteristics, developed over hundreds of millions of years, which have allowed them to adapt and survive since the beginning of life on earth. This is particularly true for human beings, so weak and frail before the occurrence of organized civilizations. The importance of effective adaptation abilities was recognized very long ago by the Chinese and Greek physicians, although it was not before the nineteenth and twentieth centuries that the real importance of the stress response with its complex multisystem mechanisms was discovered. Claude Bernard recognized the importance of a constant *milieu intérieur*, insuring the function of body cells in a changing environment via adaptive mechanisms in the vital organs. Later, Walther Cannon developed further the concept of *homeostasis*, leading in case of failure of the homeostatic mechanisms to disequilibrium and illness.

In 1936, Hans Selye published the historical Letter to the Editor of Nature, “A syndrome produced by diverse nocuous agents,” describing stress as the consequence of an inadequate response to harmful physical and psychological agents. The stress response was originally believed to be mainly related to the neuroendocrine system activation, but Selye later realized that nearly all systems were involved and the concept of a multisystem general response was developed.

Although the clinical relevance of such adaptive mechanisms in trauma and acute conditions was long recognized, their long-term effects on the mood and behavior were not acknowledged before the second part of the twentieth century: the post-traumatic stress disorder was only included in 1980 in the third edition of the DSM of the American Psychiatric Association. This underlines the central role of the brain in the response to stress, as a regulation but also as a target organ, when its vulnerability to emotional and psychological challenges, exceeds its resilience capacity and induces undesirable emotional and behavioral symptoms.

In the field of stress response, the critical care environment is unique, since it gathers individuals with different kinds of stress: patients with life-threatening conditions, health-care professionals, mainly physicians and nurses, as well as family members. In the early phase of critical illness, the patients are submitted to acute challenges such as hemorrhage, ischemia, hypoxia, sepsis and pain, as well as psychological or emotional threats. The initial adaptation to the critical illness is mainly related to the multiple autonomic, endocrine, tissue, and immune mediators; it promotes survival and recovery. However, when the critical illness is prolonged, inadequate regulation of this response may occur, inducing damaging effects, such as depressed immunity, metabolic dysfunctions, and malnutrition. At longer term, patients with prolonged or complicated stay are at risk to develop post-traumatic stress disorder, mainly characterized by psychological, emotional, and behavioral symptoms.

The family is submitted to intense psycho-emotional stress, leading to adverse psychological responses, the so-called post-intensive care syndrome-family. The latter is mainly characterized by insomnia, anxiety, depressive symptoms, inability to perform the grief, and difficulty or inability to work. Surprisingly, the clinical importance of post-intensive care syndrome-family was only discovered about 20 year ago, despite quite a high prevalence: about a third of family members are affected by PTSD symptoms in both pediatric and adult ICUs. The process of decision making involving families and proxies plays a critical role, particularly the methods of communication and inclusion. Various strategies of communication with family members have shown to be associated with decreased anxiety, improved resilience, and coping.

In addition to emotional, psychological stress, the health-care givers are submitted to work-related stress, promoting burnout symptoms, as emotional and physical exhaustion, inability to work, depersonalization, and depression. The prevalence of PTSD in ICU nurses is particularly high, due to the daily contact with suffering, uncertainty of therapy, and death. This is also the case of health-care professionals working in emergency and mental health care, which in addition are often submitted to violence and physical assaults. The occurrence and severity of work-related stress is affected by several factors, the type, nature, and severity of the stressor, the presence of a team support, and the quality of professional training, as well as by individual factors, such as the individual personality, mental health, and social-family support.

The publication of *The Stress Response of Critical Illness: Metabolic and Hormonal Aspects* by Jean-Charles Preiser and more than 20 top-level scientists must be highlighted, since it constitutes a remarkable high level and original contribution. This book comes at a right stage, as a large body of recent information has brought new insights on the metabolic and endocrine aspects of the stress response during the last decade, such as the corticoadrenal response in sepsis, the regulation of blood glucose, and substrate metabolism in the settings of critical care.

Contents

1	Introduction	1
	Jean-Charles Preiser	
Part I Metabolic Changes		
2	Successive Phases of the Metabolic Response to Stress	5
	Jean-Charles Preiser, Carole Ichai, and A.B. Johan Groeneveld	
3	Bioenergetics of the Stress Response	19
	Christophe Faisy	
4	Mitochondrial Adaptation and Hibernation	27
	Jonathan Grip, Nicolas Tardif, and Olav Rooyackers	
5	Anabolic Resistance	45
	Jean-Paul Thissen	
6	Use of Lipids as Energy Substrates	61
	Philip C. Calder and Pierre Singer	
7	The Stress Response of Critical Illness: Metabolic and Hormonal Aspects	75
	Luc Tappy	
8	Stress Hyperglycemia	89
	Jean-Charles Preiser, Aurélie Thooft, and Rafael Machado Tironi	
9	Protein Metabolism	95
	Åke Norberg, Felix Liebau, and Jan Wernerman	
10	Micronutrients	107
	Mette M. Berger	

Part II Hormonal Regulation

- 11 Thyroidal Changes During Critical Illness** 125
Lies Langouche and Greet Van den Berghe
- 12 Stress Response: Adrenal Function** 137
Paul E. Marik
- 13 Enterohormones and the Response to Critical Illness** 153
Mark P. Plummer, Annika Reintam Blaser,
and Adam M. Deane
- 14 Adipokines in Critical Illness** 169
Katherine Robinson, John Prins, and Bala Venkatesh

Part III Particular Clinical Situations

- 15 Severe Undernutrition** 187
Pierre Singer and Jonathan Cohen
- 16 The Stress Response after Traumatic Brain Injury:
Metabolic and Hormonal Aspects** 197
Hervé Quintard, Carole Ichai, and Jean-Francois Payen
- 17 Sepsis and Multiple Organ Failure** 207
Jean-Charles Preiser and Vincent Fraipont
- 18 The Stress Response of Critical Illness: Metabolic
and Hormonal Aspects, Hormonal Regulation,
Particular Clinical Situations “Morbid Obesity”** 217
Moise Coeffier and Fabienne Tamion
- 19 Hypermetabolic Response to Burn Injury** 227
Abdikarim Abdullahi, David Patsouris, Sheila R. Costford,
and Marc G. Jeschke

Chapter 1

Introduction

Jean-Charles Preiser

The important thing in science is not so much to obtain new facts as to discover new ways of thinking about them. William Henry Bragg

One of the most challenging tasks on earth is to bring people together and to build bridges to promote the cross-fertilization of knowledge!

In the field of care of the critically ill, this statement can be translated into the integration of new discoveries or major advances in the understanding of physiology into unbiased evaluations of new therapeutic strategies adapted from novel experimental findings. Conversely, basic science should be open to clinical data and able to understand and to integrate clinical concerns into research questions. This book aims to bridge the new knowledge gathered in experimental research with new clinical results.

Historically, the “*stress response*” was quoted by the Canadian physiologist Hans Selye, who discovered the mechanisms of the “fly or fight response” designed to restore the homeostasis needed for an independent life, as described by Claude Bernard. Some of the basic and adaptive mechanisms were preserved over the evolution, in keeping with the Darwinian theory of evolution.

Now, the changes and improvements in the practice of medicine allow patients to survive critical illness, allow surgeons to perform risky interventions, allow anesthesiologists to sedate very weak patients, etc. The support of the vital functions prolongs the lifespan of a critically ill thanks to improvements in pharmacological agents, in the technology of ventilators, renal replacement therapies, and extracorporeal membrane oxygenators. However, the metabolic and functional consequences of the critical illness can last weeks or months, representing a major burden for the society and cannot be supported by any dedicated device. Only a few drugs

J.-C. Preiser, MD, PhD

Department of Intensive Care, Erasme University Hospital, Université Libre de Bruxelles,
808 route de Lennik, Brussels 1070, Belgium

e-mail: Jean-Charles.Preiser@erasme.ulb.ac.be

© Springer International Publishing Switzerland 2016

J.-C. Preiser (ed.), *The Stress Response of Critical Illness:*

Metabolic and Hormonal Aspects, DOI 10.1007/978-3-319-27687-8_1

and nutritional formulas can slightly influence metabolism. Therefore, a detailed knowledge of the intimate mechanisms of the stress response is warranted to help clinicians to support the adaptive and desirable metabolic responses while trying to minimize the maladaptive ones.

In this book, several world-leading experts accepted to share their knowledge and to summarize the current understanding of the metabolic response to stress, including the cellular and subcellular mechanisms, the use of macro- and micronutrient as energy substrates during catabolism or anabolic resistance typically associated with critical illness. The roles and patterns of endocrine mediators will be discussed in detail. The final section will address difficult clinical situations, as examples of how the new findings can be translated into daily practice.

Specifically, in the first section, the bioenergetics of the stress response has been revisited in detail by C Faisy, using the concept of stress as a challenge to the equilibrium at each level of the body. The successive phases of the metabolic response to stress were being reviewed in a temporal and clinically relevant sequence. Novel insights as hibernation and mitochondrial mechanisms of adaptation have been updated by J Grip, N Tardif, and O Rooyackers, while the development of anabolic resistance has been reviewed by JP Thissen. The ensuing alterations in the use of lipids, carbohydrates, and protein metabolism were updated by Ph Calder, P Singer, L Tappy, A Norberg, F Liebau, and J Wernerman, whereas the current roles of micronutrients have been reviewed by MM Berger. A Thoof, R Machado, and myself summarized the related issue of stress hyperglycemia.

In Part 2, the current understanding of the functional changes of hormonal systems has been described by L Langouche and G Van den Berghe for the thyroid axis and P Marik for the adrenal system. The roles and relevance of new important endocrine mediators, the enterohormones released from the gastrointestinal tract, and the adipokines released from fat tissue have been reviewed by M Plummer, A Reintam, A Deane, K Robinson, J Prins, and B Venkatesh.

In Part 3, the clinical views and attitudes in situations accompanied by challenging metabolic alterations have been addressed. The issue of severe undernutrition was revisited by P Singer and J Cohen; the specificities of traumatic brain injury by H Quintard, C Ichai, and JF Payen; the particular aspects of sepsis and organ failures by V Fraipont and myself; morbid obesity by M Coeffier and F Tamion; and the issues related to burn injury by A Abdullahi, D Patsouris, SR. Costford, and MG Jeschke.

I would like to thank wholeheartedly each one of the authors who brought his own contribution and his personal stone to the huge enterprise of understanding the metabolic response to stress, an indispensable step to improve the quality of care and the quality of the lives of the survivors of critical illness.

Part I
Metabolic Changes

Chapter 2

Successive Phases of the Metabolic Response to Stress

Jean-Charles Preiser, Carole Ichai, and A.B. Johan Groeneveld

Abstract The metabolic response to stress have been selected as an adaptive response to survive critical illness. Several mechanisms well preserved over the evolution, including the stimulation of the sympathetic nervous system, the release of pituitary hormones, a peripheral resistance to the effects of these and other anabolic factors are triggered to increase the provision of energy substrates to the vital tissues. After an acute insult, alternative substrates are used as a result of the loss of control of energy substrate utilization. The clinical consequences of the metabolic response to stress include sequential changes in energy expenditure, stress hyperglycemia, changes in body composition, psychological and behavioral problems. The loss of muscle proteins and function is a major long-term consequence of stress metabolism. Specific therapeutic interventions, including hormone supplementation, enhanced protein intake and early mobilization are investigated.

2.1 Introduction

The understanding and knowledge of metabolic response to critical illness has dramatically changed during the last decade, following several important discoveries in line with the findings of pioneering scientists of the nineteenth and twentieth

J.-C. Preiser, MD, PhD (✉)

Department of Intensive Care, Erasme University Hospital, Free University of Brussels,
808 route de Lennik, Brussels 1070, Belgium
e-mail: Jean-Charles.Preiser@erasme.ulb.ac.be

C. Ichai, MD, PhD

Department of Anesthesiology and Intensive Care, Hôpital Saint-Roch, University of Nice,
Nice, France

A.B.J. Groeneveld, MD, PhD

Department of Intensive Care, Erasmus Medical Center, Rotterdam, The Netherlands

century. In his theory of the evolution, Charles Darwin reported that “It is not the strongest or the most intelligent that survives. It is the most adaptable to change.” This statement is particularly relevant after any life-threatening injury triggering a “critical illness,” when survival in a hostile environment strongly relies on the ability to mount an appropriate adaptive response. In terms of the metabolic response to stress, the principle of homeostasis of Claude Bernard (“The constancy of the internal environment is the condition for a free and independent life”) is highly relevant to the critically ill whose homeostasis must be restored as rapidly as possible to survive the injury. The mechanisms allowing the maintenance of homeostasis, vital functions, and ultimately survival in a hostile environment have been unraveled by Hans Selye, who described the “fight or flight” response, “a nonspecific response to a wide variety of stimuli.” Sir David Cuthbertson described several phases of the metabolic response over time, including the ebb phase and the flow phase. A third sequence, the chronic phase, preceding recovery, was more recently suggested and is probably relevant to the post-injury phase frequently encountered in modern intensive care [1, 2]. The mechanisms of these successive adaptive changes mounted to survive a stress are increasingly understood and are now gathered into a general theory.

2.2 Pathophysiological Mechanisms

The metabolic response to stress involves a neuroendocrine and an inflammatory/immune component. Recent data suggest that hormones released from the adipose tissue and from the gastrointestinal tract can play an important role as well (Fig. 2.1).

The neuroendocrine component is triggered in a region located near the hypothalamus, paraventricular nucleus/locus coeruleus. When a stressor is detected and signaled to the central nervous system, a prototypical response will be triggered, resulting in the activation of the sympathetic nervous system (SNS), the hypothalamic-pituitary axis, and later by behavioral changes. Many different stressors can be sensed and transmitted; for instance, a peripheral tissular injury induced by a trauma will activate afferent nerves, hypoxemia or hypercapnia will trigger chemoreceptors, hypovolemia will activate baroreceptors, and inflammatory mediators will change the phenotype of microglial cells.

The SNS is involved in the fast control of most of the body’s internal organs, via the activation of adrenergic receptors. After any stress, an immediate release of norepinephrine occurs from the postganglionic neuron in response to the stimulation of its nicotinic receptors by acetylcholine released from the preganglionic neurons [3]. The adrenal medulla is a functional sympathetic ganglion, where chromaffin cells release norepinephrine and epinephrine into the bloodstream upon stimulation by the preganglionic neuron.

The activation of the hypothalamus-pituitary axis results in the release of adrenocorticotrophic hormone, thyroid-stimulating hormone, growth hormone, and follicle-stimulating and follicle-luteinizing hormones by the anterior pituitary gland. The circulating levels of hormones released from peripheral glands in response to these




Fig. 2.1 Different levels of the metabolic response to stress. Once a stressor has been sensed, systems/organs are activated (first level). Mediators are released upon activation (second level). Physiological and phenotypical changes are triggered (third level)