



SHERLOCK'S DISEASES OF THE LIVER AND BILIARY SYSTEM

13TH EDITION

Edited by

James Dooley | Anna Lok

Guadalupe Garcia-Tsao | Massimo Pinzani

WILEY Blackwell

SHERLOCK'S DISEASES OF THE LIVER AND BILIARY SYSTEM

About the Companion Website

This book is accompanied by a companion website:

www.wiley.com/go/sherlock/liver13e

The website includes:

- PowerPoints of all figures from the book for downloading

Sherlock's Diseases of the Liver and Biliary System

Thirteenth Edition

Edited By

James S. Dooley

*University College London Institute for Liver and Digestive Health and
Sheila Sherlock Liver Centre
Royal Free London NHS Foundation Trust
London
UK*

Anna S. F. Lok

*Division of Gastroenterology and Hepatology
Michigan Medicine, University of Michigan
Ann Arbor, MI
USA*

Guadalupe Garcia-Tsao

*Section of Digestive Diseases
Yale University School of Medicine
New Haven, CT;
Veterans Affairs Connecticut Healthcare System
West Haven, CT
USA*

Massimo Pinzani

*University College London Institute for Liver and Digestive Health and
Sheila Sherlock Liver Centre
Royal Free London NHS Foundation Trust
London
UK*

WILEY Blackwell

This edition first published 2018.
© 2018 John Wiley & Sons Ltd

Edition History

Blackwell Publishing Ltd (1e, 1955; 2e, 1958; 3e, 1963; 4e, 1968; 5e, 1975; 6e, 1981; 7e, 1985; 8e, 1989; 9e, 1993; 10e, 1997; 11e, 2002; 12e, 2011)

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of James S. Dooley, Anna S. F. Lok, Guadalupe Garcia-Tsao, Massimo Pinzani to be identified as the authors of the editorial material in this work has been asserted in accordance with law.

Registered Office(s)

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA
John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Office

9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Dooley, James (James S.), editor. | Lok, Anna S. E., editor. | Garcia-Tsao, Guadalupe, editor. | Pinzani, Massimo, editor.
Title: Sherlock's diseases of the liver and biliary system / edited by James S. Dooley, Anna S.F. Lok, Guadalupe Garcia-Tsao, Massimo Pinzani.
Other titles: Diseases of the liver and biliary system
Description: 13th edition. | Hoboken, NJ : Wiley, 2018. | Preceded by Sherlock's diseases of the liver and biliary system / edited by James S. Dooley ... [et al.]. 12th ed. 2011. | Includes bibliographical references and index. |
Identifiers: LCCN 2017057075 (print) | LCCN 2017059204 (ebook) | ISBN 9781119237563 (pdf) | ISBN 9781119237648 (epub) | ISBN 9781119237549 (cloth)
Subjects: | MESH: Liver Diseases | Biliary Tract Diseases
Classification: LCC RC845 (ebook) | LCC RC845 (print) | NLM WI 700 | DDC 616.3/6--dc23
LC record available at <https://lccn.loc.gov/2017057075>

Cover image: © Nerthuz/Shutterstock
Cover design by Wiley

Set in 10/12pt Warnock by SPi Global, Pondicherry, India

Contents

- List of Contributors** *xi*
Preface to the Thirteenth Edition *xv*
Preface to the First Edition *xvi*
- 1 Anatomy and Function** *1*
Jay H. Lefkowitz
 Development of the liver and bile ducts *1*
 Anatomy of the liver *1*
 Functional liver anatomy: sectors and segments *3*
 Anatomical abnormalities of the liver *3*
 Anatomy of the biliary tract *4*
 Surface marking *5*
 Methods of examination *6*
 Microanatomy of the liver *7*
 Hepatic ultrastructure (electron microscopy) and organelle functions *9*
 Functional heterogeneity of the liver *13*
 Dynamics of the hepatic microenvironment in physiology and disease *14*
 Hepatocyte death and regeneration *15*
 References *16*
- 2 Liver Function in Health and Disease: Clinical Application of Liver Tests** *20*
George Mells and Graeme Alexander
 Bilirubin metabolism *21*
 Bile acids *23*
 Lipid and lipoprotein metabolism *26*
 Amino acid metabolism *28*
 Plasma proteins *29*
 Carbohydrate metabolism *31*
 Markers of hepatocellular injury: the serum transaminases *32*
 Markers of cholestasis: alkaline phosphatase (ALP) and gamma-glutamyl transferase (GGT) *33*
 Haematology in liver disease *34*
 Effects of ageing on the liver *35*
 References *36*
- 3 Biopsy of the Liver** *39*
David Patch and Tu Vinh Luong
 Selection and preparation of the patient *39*
 Techniques *40*
 Risks and complications *43*
 Sampling variability *45*
 Naked-eye appearances *46*
 Preparation of the specimen *46*
 Interpretation: a stepwise diagnostic approach *46*
 Indications *48*
 Special methods *49*
 References *50*
- 4 Coagulation in Cirrhosis** *53*
Nicolas M. Intagliata and Stephen H. Caldwell
 Introduction *53*
 Normal coagulation pathways: a hepatologist's perspective *54*
 The coagulation system in cirrhosis *56*
 Bleeding and thrombosis in cirrhosis *57*
 Clinical laboratory tests of the coagulation system in cirrhosis *58*
 Conclusion *60*
 References *60*
- 5 Acute Liver Failure** *62*
Shannan R. Tujios and William M. Lee
 Definition *62*
 Epidemiology and aetiologies *63*
 Clinical features *66*
 Initial investigations *67*
 Complications and management of acute liver failure *68*
 Specific therapies *73*
 Prognosis *75*
 Liver transplantation *76*
 Conclusion *78*
 References *78*

- 6 Hepatic Fibrogenesis 82**
Meena B. Bansal and Scott L. Friedman
 Introduction 82
 Natural history of hepatic fibrosis 82
 Cellular and molecular features of hepatic fibrosis 83
 Clinical aspects of hepatic fibrosis 89
 Emerging antifibrotic targets and strategies 89
 References 90
- 7 Non-invasive Assessment of Fibrosis and Cirrhosis 93**
Avik Majumdar and Massimo Pinzani
 Introduction 93
 The use of invasive and non-invasive tests 93
 Non-invasive tests: specifics 95
 Conclusions 102
 References 103
- 8 Hepatic Cirrhosis 107**
P. Aiden McCormick and Rajiv Jalan
 Definition 107
 Causes of cirrhosis 107
 Anatomical diagnosis 108
 Reversible cirrhosis 110
 Clinical cirrhosis: compensated versus decompensated 110
 Prognosis (Child–Pugh score, MELD, UKELD) 111
 Clinical and pathological associations 112
 Management 119
 Acute-on-chronic liver failure 120
 References 123
- 9 Ascites 127**
Guadalupe Garcia-Tsao
 Mechanisms of ascites formation 127
 Clinical features 130
 Differential diagnosis 132
 Spontaneous bacterial peritonitis 132
 Treatment of cirrhotic ascites 134
 Hyponatraemia 138
 Refractory ascites 139
 Hepatorenal syndrome 141
 Prognosis 144
 References 145
- 10 Hepatic Encephalopathy in Patients with Cirrhosis 151**
Marsha Y. Morgan
 Clinical Features 151
 Classification 153
 Prevalence and consequences 154
 Diagnosis 154
 Diagnostic comorbidities, confounders, and alternatives 160
 Pathogenesis 161
 Management 167
 Prevention 174
 References 174
- 11 Portal Hypertension in Cirrhosis 180**
Jaime Bosch and Annalisa Berzigotti
 Introduction 180
 Pathophysiology and rational basis of therapy 185
 Evaluation and diagnosis 188
 Natural history and prognosis 194
 Management 196
 Treatment of portal hypertension according to clinical scenarios 200
 References 205
- 12 Vascular Disorders of the Liver and Extrahepatic Portal Hypertension 209**
Dominique-Charles Valla
 Hepatic artery occlusion 209
 Aneurysms of the hepatic artery 210
 Hepatic arterioportal fistula 211
 Hepatic vascular malformations in hereditary haemorrhagic telangiectasia 212
 Congenital portosystemic shunts – Abernethy malformation 213
 Budd–Chiari syndrome – hepatic venous outflow tract obstruction 214
 Extrahepatic portal vein obstruction – portal vein thrombosis and portal cavernoma in the absence of cirrhosis 217
 Portal vein thrombosis in patients with cirrhosis 220
 Idiopathic non-cirrhotic intrahepatic portal hypertension 221
 Hypoxic hepatitis 223
 Congestive cardiac hepatopathy 223
 Non-obstructive sinusoidal dilation (NOSD) and peliosis 225
 References 226
- 13 Jaundice and Cholestasis 231**
Peter L. M. Jansen
 Introduction 231
 Mechanics of bile formation 233
 Syndrome of cholestasis 238
 Causes of isolated hyperbilirubinaemia 239
 Causes of cholestatic and hepatocellular jaundice 242

- Consequences of cholestasis and their management 244
Investigation of the jaundiced patient 247
Decisions to be made in the jaundiced patient 250
Management of cholestatic disorders 251
References 251
- 14 Gallstones and Benign Biliary Disease 256**
James S. Dooley, Kurinchi S. Gurusamy, and Brian R. Davidson
Introduction 256
Imaging the gallbladder and biliary tract 256
Gallstones 259
Symptoms and complications of gallstones 264
Cholecystectomy 266
Complicated acute gallbladder disease 268
Percutaneous cholecystostomy 269
Asymptomatic gallbladder stones 269
Non-surgical treatment of gallstones in the gallbladder 269
Common bile duct stones 270
Acute gallstone pancreatitis 272
Large common duct stones 272
Mirizzi syndrome 273
Intrahepatic gallstones 274
Haemobilia 274
Functional gallbladder and sphincter of Oddi disorders 274
Other gallbladder pathologies 276
Relationships to malignant change 279
Benign biliary strictures 279
Anastomotic strictures following biliary surgery 282
IgG4-related sclerosing cholangitis 283
Chronic pancreatitis 283
References 284
- 15 Malignant Biliary Diseases 294**
Rahul S. Koti and John Bridgewater
Carcinoma of the gallbladder 294
Carcinoma of the bile duct (cholangiocarcinoma) 296
Other biliary malignancies 301
Metastases at the hilum 301
Ampullary and periampullary carcinomas 301
Conclusion 305
References 305
- 16 Fibropolycystic Liver Diseases and Congenital Biliary Abnormalities 308**
Nedim Hadžić and Mario Strazzabosco
Overview 308
Polycystic liver disease 309
Fibropolycystic diseases 313
Autosomal recessive polycystic kidney disease 314
Congenital hepatic fibrosis 314
Caroli disease 316
Microhamartomas (von Meyenberg complexes) 318
Choledochal cysts 318
Solitary non-parasitic liver cyst 320
Congenital anomalies of the biliary tract 321
References 324
- 17 Primary Biliary Cholangitis 328**
David E. J. Jones
Clinical features 328
Diagnosis 329
Epidemiology 333
Aetiology and pathogenesis 334
Management 335
Prognosis 337
References 338
- 18 Sclerosing Cholangitis 341**
Tom Hemming Karlsen and Kirsten Muri Boberg
Introduction 341
Primary sclerosing cholangitis 341
Secondary sclerosing cholangitis 350
Sclerosing cholangitis in systemic inflammatory diseases 351
References 351
- 19 Autoimmune Hepatitis and Overlap Syndromes 355**
Ashnila Janmohamed and Gideon M. Hirschfield
Introduction 355
Disease overview 356
Biological determinants of disease 357
Disease presentation 359
Laboratory features 361
Imaging 363
Liver biopsy and histological features 363
Differential diagnosis 365
Diagnostic dilemmas 366
Making a diagnosis in practice 367
Management strategies 368
Pretreatment and on-treatment considerations 371
Treatment challenges and alternative agents 371
Pregnancy and autoimmune hepatitis 372
The elderly and autoimmune hepatitis 372
Childhood-onset autoimmune hepatitis 373
Autoimmune hepatitis and liver transplantation 373
Overlap syndromes 374
Conclusion 377
References 377

20 Enterically Transmitted Viral Hepatitis: Hepatitis A and Hepatitis E 380*Rinjal D. Brahmabhatt and Anna S. F. Lok*

General features of enterically transmitted viral hepatitis 380

Hepatitis A virus 385

Hepatitis E virus 389

References 391

21 Hepatitis B 395*Anna S. F. Lok*

Introduction 395

Hepatitis B virus 395

Immune response and mechanisms of hepatic injury 398

Epidemiology 399

Prevention 400

Diagnosis 402

Clinical manifestations 404

Natural history 405

Treatment 408

HBV and HCV coinfection 414

HBV and HDV coinfection 414

HBV and HIV coinfection 414

References 415

22 Hepatitis D 421*Patrizia Farci and Grazia Anna Niro*

History 421

Hepatitis D virus 421

Epidemiology 422

Pathogenesis 425

Modes of infection and clinical course 425

Diagnosis 428

Treatment 429

Prevention 432

References 432

23 Hepatitis C 436*Geoffrey Dusheiko*

Introduction 436

Epidemiology 436

Virology 438

Pathology and pathogenesis 439

Diagnostic tests for hepatitis C 440

Acute hepatitis C 441

Chronic hepatitis C 441

References 459

24 Drug-Induced Liver Injury 468*Frank W. DiPaola and Robert J. Fontana*

Introduction 468

Epidemiology 468

Complications of DILI 469

Classification of hepatotoxicity 469

Drug metabolism and pharmacokinetics 469

Hepatic drug metabolism 471

Molecular mechanisms in drug-induced liver injury 472

Non-genetic risk factors for DILI 473

Diagnosis of DILI 474

Medical management 478

Pharmacogenetic risk factors 478

Potential immunological mechanisms in idiosyncratic DILI 479

Liver injury from specific drugs 479

References 486

25 Alcohol and the Liver 494*Stephen Stewart and Ewan Forrest*

Introduction 494

Alcohol metabolism 494

Pathogenesis 496

Susceptibility 497

Histological features 498

Clinical features 501

Clinical syndromes 503

Prognosis 504

Treatment 506

Conclusions 507

References 508

26 Iron Overload States 511*Paul Adams and Heinz Zoller*

Normal iron physiology 511

Iron overload and liver damage 515

Genetic haemochromatosis 516

Other iron storage diseases 521

References 523

27 Wilson Disease 526*Eve A. Roberts and Karl Heinz Weiss*

Molecular genetics: pathogenesis 527

Pathology 528

Clinical picture 529

Laboratory tests 532

Genetic strategies 533

Diagnostic difficulties 533

Treatment 533

Prognosis 536

Non-Wilsonian copper-related cirrhosis 536

References 536

28 Non-Alcoholic Fatty Liver Disease 540*Timothy Hardy and Christopher P. Day*

Introduction 540

Further definitions, terminology, and diagnosis 541

Liver biopsy, classification of NAFLD, and non-invasive markers of NASH and fibrosis 541

- Clinical features 543
 Laboratory testing 544
 Epidemiology 545
 Ethnic variation in NAFLD 545
 Pathogenesis of NASH 545
 Natural history of NAFLD 550
 NAFLD and hepatocellular carcinoma (HCC) 551
 Therapy for non-alcoholic fatty liver disease 552
 Other forms of NAFLD 554
 References 555
- 29 Nutrition and Chronic Liver Disease 561**
Manuela Merli
 Introduction 561
 Epidemiology and general characteristics 562
 Causes of malnutrition 562
 Consequences of malnutrition 564
 Diagnosis and assessment 565
 Treatment and management 568
 References 570
- 30 Pregnancy and the Liver 572**
Rachel H. Westbrook and Catherine Williamson
 Introduction 572
 Normal physiology in pregnancy 572
 Pregnancy-related liver diseases 573
 Pre-existing liver diseases and pregnancy 578
 Liver transplantation and pregnancy 580
 Liver disease coincidentally arising with pregnancy 581
 Conclusion 582
 References 582
- 31 The Liver in the Neonate, in Infancy, and Childhood 588**
Susan M. Siew and Deirdre A. Kelly
 Investigation of liver disease in children 588
 Neonatal jaundice 589
 Neonatal unconjugated hyperbilirubinaemia 589
 Neonatal liver disease (conjugated hyperbilirubinaemia) 591
 Neonatal hepatitis 594
 Inherited disease in the neonate 596
 Genetic cholestatic syndromes 598
 Structural abnormalities: biliary atresia and choledochal cyst 600
 Acute liver failure in infancy 602
 Liver disease in older children 605
 Metabolic disease in older children 607
 Cirrhosis and portal hypertension 613
 Liver transplantation 613
 Tumours of the liver 614
 References 615
- 32 The Liver in Systemic Diseases 622**
James S. Dooley and Christopher McNamara
 Collagen-vascular and autoimmune disorders 622
 Hepatic granulomas 624
 Sarcoidosis 626
 The liver in endocrine disorders 628
 Amyloidosis 629
 Porphyrias 632
 The liver in haemolytic anaemias 634
 The liver in myelo- and lymphoproliferative disease 638
 Bone marrow transplantation 639
 Lymphoma 640
 Extramedullary haemopoiesis 642
 Rare haematological disorders that may involve the liver 643
 Lipid storage diseases 643
 Non-metastatic complications of malignancy 646
 References 646
- 33 The Liver in Infections 652**
Sanjay Bhagani and Ian Cropley
 Introduction 652
 Jaundice of infections 652
 Pyogenic liver abscess 652
 Hepatic amoebiasis 655
 Tuberculosis of the liver 657
 Hepatic actinomycosis 659
 Syphilis of the liver 659
 Perihepatitis 660
 Leptospirosis 660
 Relapsing fever 663
 Lyme disease 663
 Rickettsial infections 663
 Fungal infections 664
 Schistosomiasis (bilharzia) 665
 Malaria 667
 Kala-azar (visceral leishmaniasis) 668
 Echinococcosis (hydatid disease) 668
 Ascariasis 673
 Strongyloides stercoralis 674
 Trichinosis 674
 Toxocara canis (visceral larva migrans) 674
 Liver flukes 675
 References 676
- 34 Imaging of the Liver and Diagnostic Approach of Space-Occupying Lesions 682**
Neil H. Davies and Dominic Yu
 Ultrasound 682
 Computed tomography 683
 Magnetic resonance imaging 685
 Radioisotope scanning 688
 Positron emission tomography 691

MR spectroscopy 691
Conclusions and choice of imaging technique 691
References 691

35 Benign Liver Tumours 693

Ian R. Wanless

Diagnosis of focal liver lesions 693
Hepatocellular lesions 693
Biliary and cystic lesions 699
Mesenchymal tumours 700
References 701

36 Primary Malignant Neoplasms of the Liver 705

Adam Doyle and Morris Sherman

Hepatocellular carcinoma 705
Intrahepatic cholangiocarcinoma 718
Other malignant neoplasms of the liver 721
Other sarcomas 722
References 722

37 Hepatic Transplantation 730

Lindsay Y. King and Carl L. Berg

Selection of patients 730
Candidates 732

Absolute and relative contraindications 737
General preparation of the patient 738
Donor selection and operation 738
The recipient operation 739
Immunosuppression 742
Postoperative course 743
Post-transplantation complications 744
Conclusion 751
References 752

38 Hepatic Transplantation and HBV, HCV, and HIV

Infections 758

Norah A. Terrault

Introduction 758
Hepatitis B and liver transplantation 759
Hepatitis C and liver transplantation 763
HIV and liver transplantation 770
References 773

Index 781

List of Contributors

Paul Adams MD

Professor of Medicine
Chief of Gastroenterology
University Hospital
University of Western Ontario
London, ON, Canada

Graeme Alexander MA, MD, FRCP

UCL Professor
Institute for Liver and Digestive Health
UCL Medical School
London, UK

Meena B. Bansal MD

Associate Professor of Medicine
Division of Liver Diseases
Icahn School of Medicine at Mount Sinai
New York, NY, USA

Carl L. Berg MD

Professor of Medicine
Medical Director of Abdominal Transplantation
Duke University Health System
Durham, NC, USA

Annalisa Berzigotti MD, PhD

Swiss Liver Center, Hepatology
University Clinic for Visceral Surgery and Medicine
Inselspital
University of Bern
Bern, Switzerland

Sanjay Bhagani BSc, MB BCh, FRCP

Consultant Physician and Honorary Senior Lecturer
Department of Infectious Diseases/HIV Medicine
Royal Free Hospital
Research Department of Infection, UCL
London, UK

Kirsten Muri Boberg MD, PhD

Professor of Hepatology
Department of Transplantation Medicine
Oslo University Hospital Rikshospitalet
Oslo, Norway

Jaime Bosch MD, PhD, FRCP, FAASLD

Hepatic Hemodynamic Laboratory, Liver Unit,
Hospital Clinic, IDIBAPS, University of Barcelona
Barcelona, Spain;
CIBERehd, Instituto de Salud Carlos III, Madrid, Spain;
Swiss Liver Center, Hepatology, University Clinic for
Visceral Surgery and Medicine, Inselspital, University
of Bern
Bern, Switzerland

Rinjal D. Brahmabhatt MD

Division of Gastroenterology and Hepatology
Michigan Medicine, University of Michigan
Ann Arbor, MI, USA

John Bridgewater MA, FRCP, PhD, MB BS

Professor and Consultant in Medical Oncology
University College London Medical School and Cancer
Institute
London, UK

Stephen H. Caldwell MD

Professor, Director of Hepatology
Division of Gastroenterology and Hepatology
University of Virginia
Charlottesville, VA, USA

Ian Cropley MA, MB BS, FRCP, DTM&H

Consultant Physician and Honorary Senior Lecturer
Department of Infectious Diseases/HIV Medicine
Royal Free Hospital
London, UK

Brian R. Davidson MB BCh, MD, FRCS (Glas), FRCS (Eng)

Consultant HPB and Liver Transplant Surgeon
HPB and Liver Transplant Unit
Royal Free Hospital, London
Professor of Surgery
Head of HBP and Liver Transplant Research
University College London Medical School, Royal Free
Campus,
London, UK

Neil H. Davies MB BS, FRCS, FRCR
Consultant Interventional Radiologist
Department of Radiology
Royal Free Hospital
London, UK

Christopher P. Day FMedSci
Professor of Liver Medicine
Institute of Cellular Medicine, Newcastle University
Newcastle upon Tyne, UK

Frank W. DiPaola MD
Department of Pediatrics
University of Michigan Medical School
Ann Arbor, MI, USA

James S. Dooley MD, FRCP
Emeritus Reader in Medicine, UCL Institute for Liver
and Digestive Health, and Honorary Consultant
Hepatologist, Sheila Sherlock Liver Centre,
Royal Free Hospital
London, UK

Adam Doyle MBBS, BMedSc, FRACP
Department of Gastroenterology and Hepatology
Royal Perth Hospital
Perth, WA, Australia

Geoffrey Dusheiko FCP(SA), FRCP, FRCP(Edin)
Emeritus Professor of Medicine, UCL Institute
for Liver and Digestive Health, and Kings College
Hospital
London, UK

Patrizia Farci MD
Chief, Hepatic Pathogenesis Section
Laboratory of Infectious Diseases
National Institute of Allergy and Infectious Diseases
National Institutes of Health
Bethesda, MD, USA

Robert J. Fontana, MD
Professor of Medicine
Department of Internal Medicine, University of
Michigan Medical School
Ann Arbor, MI, USA

Ewan Forrest MD, FRCPS (Glasgow), FRCP (Edin)
Consultant Hepatologist
Liver Unit, Glasgow Royal Infirmary
Glasgow, UK

Scott L. Friedman MD
Fishberg Professor of Medicine
Chief, Division of Liver Diseases
Icahn School of Medicine at Mount Sinai
New York, NY, USA

Guadalupe Garcia-Tsao MD
Professor of Medicine
Section of Digestive Diseases
Yale School of Medicine
New Haven, CT;
Veterans Affairs Connecticut Healthcare System
West Haven, CT, USA

Kurinchi S. Gurusamy MB BS, MRCS, PhD
Reader in Surgery
University College London Medical School, Royal Free
Campus, and Royal Free Hospital, London, UK

Nedim Hadžić MD, MSc, FRCPCH, FAALD
Professor and Consultant in Paediatric Hepatology
Paediatric Centre for Hepatology, Gastroenterology and
Nutrition
King's College Hospital
London, UK

Timothy Hardy MB BS, BSc (Hons)
MRC Clinical Research Fellow
Institute of Cellular Medicine, Newcastle University
Newcastle upon Tyne, UK

Gideon M. Hirschfield MA, MB BChir, PhD, FRCP
Professor and Honorary Consultant Hepatologist
Centre for Liver Research, NIHR Biomedical Research
Centre, University of Birmingham
Birmingham, UK

Nicolas M. Intagliata MD
Assistant Professor of Medicine, Gastroenterology and
Hepatology
University of Virginia
Charlottesville, VA, USA

Rajiv Jalan MBBS, MD, PhD, FRCP, FRCPE, FAASLD
Professor of Hepatology
Liver Failure Group, Institute for Liver and Digestive
Health, Division of Medicine, University College
London
London, UK

Ashnila Janmohamed BSc, MBBS, MRCP
Clinical Research Fellow in Hepatology
Centre for Liver Research, NIHR Biomedical Research
Centre, University of Birmingham
Birmingham, UK

Peter L. M. Jansen MD, PhD

Professor of Hepatology
 Department of Gastroenterology and Hepatology
 Academic Medical Center
 University of Amsterdam
 Amsterdam, The Netherlands

David E. J. Jones BM, BCh, MA, PhD, FRCP

Professor of Liver Immunology
 Medical School, Newcastle University
 Honorary Consultant Hepatologist
 Freeman Hospital
 Newcastle upon Tyne, UK

Tom Hemming Karlsen MD, PhD

Professor of Internal Medicine
 Senior Consultant of Gastroenterology and Hepatology
 Department of Transplantation Medicine
 Oslo University Hospital Rikshospitalet
 Oslo, Norway

Deirdre A. Kelly MD, FRCP, FRCPI, FRCPC

Professor of Paediatric Hepatology
 Birmingham Children's Hospital and University of
 Birmingham
 Birmingham, UK

Lindsay Y. King MD, MPH

Assistant Professor of Medicine
 Duke University Health System
 Durham, NC, USA

Rahul S. Koti MD, FRCS

Honorary Senior Lecturer in Surgery
 Royal Free Hospital and University College London
 Medical School
 London, UK

William M. Lee MD, FACP

Professor of Internal Medicine
 University of Texas Southwestern Medical Center
 Dallas, TX, USA

Jay H. Lefkowitz MD

Professor of Clinical Pathology
 College of Physicians and Surgeons, Columbia University
 New York, NY, USA

Anna S. F. Lok MBBS, MD, FRCP

Alice Lohrman Andrews Research Professor in
 Hepatology
 Director of Clinical Hepatology
 Division of Gastroenterology and Hepatology
 Michigan Medicine, University of Michigan
 Ann Arbor, MI, USA

Tu Vinh Luong MD

Consultant Histopathologist
 Department of Cellular Pathology, Royal Free Hospital;
 Honorary Senior Lecturer,
 UCL Institute for Liver and Digestive Health
 London, UK

Avik Majumdar MBBS(Hons), MPHTM, FRACP

UCL Institute for Liver and Digestive Health and
 Sheila Sherlock Liver Centre
 Royal Free Hospital, London, UK;
 AW Morrow Gastroenterology and Liver Centre
 Australian National Liver Transplant Unit
 Royal Prince Alfred Hospital,
 Sydney, NSW, Australia

Christopher McNamara FRACP, FRCPA, FRCPath

Department of Haematology
 University College London Hospitals
 London, UK

P. Aiden McCormick MD, FRCP, FRCPI, FEBGH, FAASLD

Consultant Hepatologist and Newman
 Clinical Research Professor
 St Vincent's University Hospital and
 University College Dublin
 Dublin, Ireland

George Mells MB BS, PhD, MRCP

Consultant Hepatologist
 Cambridge Liver Unit
 Addenbrooke's Hospital
 Cambridge, UK

Manuela Merli MD

Associate Professor, Gastroenterology
 Department of Clinical Medicine, Sapienza
 University of Rome
 Rome, Italy

Marsha Y. Morgan MB ChB, FRCP

Principal Research Associate and Honorary
 Consultant Physician
 UCL Institute for Liver and Digestive Health
 Division of Medicine
 University College London
 London, UK

Grazia Anna Niro, MD

Gastroenterology Unit
 IRCCS "Casa Sollievo Sofferenza" Hospital
 San Giovanni Rotondo
 Foggia, Italy

David Patch MB BS, FRCP

Consultant Hepatologist
Sheila Sherlock Liver Centre, Royal Free Hospital and
UCL Institute for Liver and Digestive Health
London, UK

Massimo Pinzani MD, PhD, FRCP, FAASLD

Professor of Medicine
UCL Institute for Liver and Digestive Health and Sheila
Sherlock Liver Centre
Royal Free Hospital
London, UK

Eve A. Roberts MD, PhD, FRCPC

Departments of Paediatrics, Medicine, and
Pharmacology and Toxicology
University of Toronto
Toronto, ON, Canada

Morris Sherman MB BCh, PhD, FRCP(C), FAASLD

Professor of Medicine
Department of Medicine,
Division of Gastroenterology
University of Toronto and
University Health Network
Toronto, ON, Canada

Susan M. Siew MBBS (Hons.), FRACP, PhD

Paediatric Gastroenterologist
Department of Gastroenterology and James Fairfax
Institute of Paediatric Nutrition
Sydney Children's Hospital Network (Westmead)
Sydney, NSW, Australia

Stephen Stewart MBChB, PhD

Consultant Hepatologist and Director of Centre for
Liver Disease
Mater Misericordiae University Hospital
Dublin, Ireland

Mario Strazzabosco MD, PhD

Professor of Medicine;
Director, Smilow Liver Cancer Program;
Deputy Director, Yale Liver Center
Yale School of Medicine
New Haven, CT, USA

Norah A. Terrault MD, MPH

Professor of Medicine
Division of Gastroenterology/Hepatology
University of California San Francisco
San Francisco, CA, USA

Shannan R. Tujios MD

Assistant Professor of Internal Medicine
Division of Digestive and Liver Diseases
University of Texas Southwestern Medical Center
Dallas, TX, USA

Dominique-Charles Valla MD

Professor of Hepatology
University Paris-Diderot and Inserm UMR-1149, Paris;
Department of Hepatology, Hôpital Beaujon, APHP
Clichy-la-Garenne, France

Ian R. Wanless MD CM, FRCPC

Professor of Pathology
Department of Pathology
Queen Elizabeth II Health Sciences Centre and
Dalhousie University
Halifax, NS, Canada

Karl Heinz Weiss MD

Head of Section of Transplant Hepatology
Department of Internal Medicine IV
University of Heidelberg
Heidelberg, Germany

Rachel H. Westbrook MBChB, MRCP, MD

Consultant Hepatologist and Honorary Senior Lecturer
Royal Free Hospital and UCL Institute for Liver and
Digestive Health
London, UK

Catherine Williamson MD, FRCP

Professor of Women's Health, Department of Women
and Children's Health
Faculty of Life Sciences & Medicine
King's College London
London, UK

Dominic Yu MB BS, MRCPI, FRCR

Consultant Radiologist
Department of Radiology
Royal Free Hospital
London, UK

Heinz Zoller MD

Associate Professor of Medicine
Consultant in Gastroenterology and Hepatology
Department of Internal Medicine 1
Medical University of Innsbruck
Innsbruck, Austria

Preface to the Thirteenth Edition

Since 2011 important new data have contributed to all areas of hepatobiliary disease, and revision of the 12th edition of *Sherlock's Diseases of the Liver and Biliary System*, published then, is timely. Many clinical challenges remain. However, advances in the understanding of pathogenesis and prognostic staging are leading to more individualized patient care and providing improved outcomes in many conditions. Studies of the basic cellular mechanisms involved are leading to the investigation of new therapeutic approaches.

A particular advance with important implications is in the treatment of hepatitis C, which can now be cured in more than 95% of patients with 8 to 16 weeks of oral antiviral therapy.

The current hepatological epidemic in the Western world is non-alcoholic fatty liver disease and much research is published and on-going to dissect and understand the mechanisms and predisposing factors related to progression to cirrhosis. New agents are being studied for their potential to prevent or reverse such progression.

The drive of clinical hepatologists and basic scientists for greater understanding is a credit to the worldwide community of those involved. International collaborations to gather data on large cohorts of patients are playing an important role. Viral hepatitis has been mentioned, but there has been much new work in cholestatic liver diseases. Molecular genetic techniques are more accessible, for example in the diagnosis of conditions such as haemochromatosis and Wilson disease. Gene therapy may be on the horizon for some genetic diseases. Genome-wide studies are becoming almost routine to dissect the predisposing factors, for example in immune-related and alcoholic liver disease. Liver imaging, already strides ahead of what was available in earlier days, has greatly reduced the need for biopsies. Liver transplantation is widely available. The selection and management of this cohort of patients is constantly being assessed and refined to attain better outcomes.

Sheila Sherlock, an enthusiastic pioneer in the development of Hepatology worldwide, would have been

excited and we think stunned by the current state of her speciality. She herself used her textbook as a resource in which to annotate the most current work and knowledge. Her goal with each edition was to provide a textbook of manageable size, critical and current.

It is with this ethos that we have approached the 13th edition with specialist contributors from across the world. We hope that this edition will stimulate interest in hepatobiliary disease among students and young doctors, as well as provide a reference source for specialist hepatologists, surgeons, pathologists, radiologists, and clinical nurse specialists. The edition has over 100 new figures and more than 1500 new references.

The format is as before – designed to preserve clarity, with the text written as far as is possible with Sheila Sherlock's style in mind.

New chapters reflect current developments and interest with particular contributions on non-invasive assessment of hepatic fibrosis and cirrhosis, coagulation in cirrhosis, and nutrition in the patient with cirrhosis.

We are most grateful to the authors for their work updating, fully revising, or providing new chapters. We thank Pri Gibbons and Deirdre Barry at Wiley Blackwell for their contributions to making this project a success. We wish also to express our particular thanks to Gill Whitley for her unflinching commitment, expertise, and support during the production of this 13th edition. We are grateful to Jane Fallows for the new artwork and for the reworking where necessary of figures from the previous edition.

Finally, this textbook is the legacy of Sheila Sherlock's exceptional life and career, and we hope that her daughters and their families will take pleasure in seeing their mother's seminal contribution to Hepatology celebrated in another edition of *Diseases of the Liver and Biliary System*.

September 2017

James S. Dooley
Anna S. F. Lok
Guadalupe Garcia-Tsao
Massimo Pinzani

Preface to the First Edition

My aim in writing this book has been to present a comprehensive and up-to-date account of diseases of the liver and biliary system, which I hope will be of value to physicians, surgeons, and pathologists and also a reference book for the clinical student. The modern literature has been reviewed with special reference to articles of general interest. Many older more specialized classical contributions have therefore inevitably been excluded.

Disorders of the liver and biliary system may be classified under the traditional concept of individual diseases. Alternatively, as I have endeavoured in this book, they may be described by the functional and morphological changes which they produce. In the clinical management of a patient with liver disease, it is important to assess the degree of disturbance of four functional and morphological components of the liver: hepatic cells, vascular system (portal vein, hepatic artery, and hepatic veins), bile ducts, and reticulo-endothelial system. The typical reaction pattern is thus sought and recognized before attempting to diagnose the causative insult. Clinical and laboratory methods of assessing each of these components are therefore considered early in the book. Descriptions of individual diseases follow as illustrative examples. It will be seen that the features of hepatocellular failure and portal hypertension are described in general terms as a foundation for subsequent discussion of virus hepatitis, nutrition liver disease, and the cirrhotoses. Similarly blood diseases and infections of the liver are included with the reticuloendothelial system, and disorders of the biliary tract follow descriptions of acute and chronic bile duct obstruction.

I would like to acknowledge my indebtedness to my teachers, the late Professor J. Henry Dible, the late Professor Sir James Learmonth and Professor Sir John McMichael, who stimulated my interest in hepatic disease, and to my colleagues at the Postgraduate Medical School and elsewhere who have generously invited me to see patients under their care. I am grateful to Dr A. G. Bearn for criticizing part of the typescript and to Dr A. Paton for his criticisms and careful proof reading. Miss D. F. Atkins gave much assistance with proof reading and

with the bibliography. Mr Per Saugman and Mrs J. M. Green of Blackwell Scientific Publications have co-operated enthusiastically in the production of this book.

The photomicrographs were taken by Mr E. V. Willmott, FRPS, and Mr C. A. P. Graham from sections prepared by Mr J. G. Griffin and the histology staff of the Postgraduate Medical School. Clinical photographs are the work of Mr C. R. Brecknell and his assistants. The black and white drawings were made by Mrs H. M. G. Wilson and Mr D. Simmonds. I am indebted to them all for their patience and skill.

The text includes part of unpublished material included in a thesis submitted in 1944 to the University of Edinburgh for the degree of MD, and part of an essay awarded the Buckston–Browne prize of the Harveian Society of London in 1953. Colleagues have allowed me to include published work of which they are jointly responsible. Dr Patricia P. Franklyn and Dr R. E. Steiner have kindly loaned me radiographs. Many authors have given me permission to reproduce illustrations and detailed acknowledgments are given in the text. I wish also to thank the editors of the following journals for permission to include illustrations: *American Journal of Medicine*, *Archives of Pathology*, *British Heart Journal*, *Circulation*, *Clinical Science*, *Edinburgh Medical Journal*, *Journal of Clinical Investigation*, *Journal of Laboratory and Clinical Investigation*, *Journal of Pathology and Bacteriology*, *Lancet*, *Postgraduate Medical Journal*, *Proceedings of the Staff Meetings of the Mayo Clinic*, *Quarterly Journal of Medicine*, *Thorax*, and also the following publishers: Butterworth's Medical Publications, J. & A. Churchill Ltd, The Josiah Macy Junior Foundation, and G. D. Searle & Co.

Finally I must thank my husband, Dr D. Geraint James, who, at considerable personal inconvenience, encouraged me to undertake the writing of this book and also criticized and rewrote most of it. He will not allow me to dedicate it to him.

SHEILA SHERLOCK
1955

Chapter 1

Anatomy and Function

Jay H. Lefkowitz

College of Physicians and Surgeons, Columbia University, New York, NY, USA

LEARNING POINTS

- The liver is derived from a foregut endodermal bud which develops in the third week of gestation and divides into two parts: hepatic and biliary.
- The Couinaud classification subdivides the liver into eight segments (segments I–IV in the left lobe, segments V–VIII in the right lobe) based on vascular and biliary anatomical landmarks.
- The lobule described by Kiernan is the most widely used unit of liver microanatomy, consisting of a hexagon-like region of liver parenchyma with a central vein as its hub and portal tracts located in the periphery of the hexagon.
- Hepatocytes are functionally heterogeneous within the lobular parenchyma, whereby centrilobular cells subserve different functions (e.g. drug metabolism) from periportal cells (e.g. bile salt-dependent bile formation).
- Uncomplicated regeneration of hepatocytes and/or bile duct epithelium usually occurs by cell division of the indigenous cells; however, when normal regenerative capacity is overwhelmed there may be activation of progenitor cells located in the region of the canals of Hering.

Development of the liver and bile ducts

The liver begins as a hollow endodermal bud from the foregut (duodenum) during the third week of gestation. The bud separates into two parts: hepatic and biliary. The *hepatic* part contains bipotential progenitor cells that differentiate into hepatocytes or ductal cells, which form the early primitive bile duct structures (bile duct plates). Differentiation is accompanied by changes in cytokeratin type within the cell [1]. Normally, this collection of rapidly

proliferating cells penetrates adjacent mesodermal tissue (the septum transversum) and is met by ingrowing capillary plexuses from the vitelline and umbilical veins, which will form the sinusoids. The connection between this proliferating mass of cells and the foregut, the *biliary* part of the endodermal bud, will form the gallbladder and extrahepatic bile ducts. The intrahepatic bile ducts begin to develop at week 8 from the flattened epithelium of the ductal plates adjacent to portal tracts [2]. Bile begins to flow at about the 12th week. Connective tissue cells of portal tracts are derived from the mesoderm of the septum transversum. Kupffer cells derive from circulating monocytes and possibly yolk sac macrophages. Hepatic stellate cells appear to be mesodermal derivatives from submesothelial cells located beneath the surface of the developing liver [3]. The fetal liver is the main site of haemopoiesis by the 12th week; this subsides in the fifth month coincident with the onset of bone marrow haemopoietic activity, so that only a few haemopoietic cells remain at birth.

Anatomy of the liver

The liver, the largest organ in the body, weighs 1200–1500g and comprises one-fiftieth of the total adult body weight. It is relatively larger in infancy, comprising one-eighteenth of the birth weight. This is mainly due to a large left lobe.

Sheltered by the ribs in the right upper quadrant, the upper border lies approximately at the level of the nipples. There are two anatomical lobes, the right being about six times the size of the left (Fig. 1.1, Fig. 1.2, Fig. 1.3). Lesser segments of the right lobe are the *caudate lobe* on the posterior surface and the *quadrate lobe* on the inferior surface. The right and left lobes are separated anteriorly by a fold of peritoneum called the falciform ligament, posteriorly by the fissure for the ligamentum venosum, and inferiorly by the fissure for the ligamentum teres.

Sherlock's Diseases of the Liver and Biliary System, Thirteenth Edition. Edited by James S. Dooley, Anna S. F. Lok, Guadalupe Garcia-Tsao and Massimo Pinzani.

© 2018 John Wiley & Sons Ltd. Published 2018 by John Wiley & Sons Ltd.

Companion website: www.wiley.com/go/sherlock/liver13e

The liver has a double blood supply. The *portal vein* brings venous blood from the intestines and spleen and the *hepatic artery*, coming from the coeliac axis, supplies the liver with arterial blood. These vessels enter the liver through a fissure, the *porta hepatis*, which lies far back on the inferior surface of the right lobe. Inside the porta, the portal vein and hepatic artery divide into branches to the right and left lobes, and the right and left hepatic bile ducts




Fig. 1.1 Anterior view of the liver.




Fig. 1.2 Posterior view of the liver.




Fig. 1.3 Inferior view of the liver.

join to form the common hepatic duct. The *hepatic nerve plexus* contains fibres from the sympathetic ganglia T7–T10, which synapse in the coeliac plexus, the right and left vagi and the right phrenic nerve. It accompanies the hepatic artery and bile ducts into their finest ramifications, even to the portal tracts and hepatic parenchyma [4].

The *ligamentum venosum*, a slender remnant of the ductus venosus of the fetus, arises from the left branch of the portal vein and fuses with the inferior vena cava at the entrance of the left hepatic vein. The *ligamentum teres*, a remnant of the umbilical vein of the fetus, runs in the free edge of the falciform ligament from the umbilicus to the inferior border of the liver and joins the left branch of the portal vein. Small veins accompanying it connect the portal vein with veins around the umbilicus. These become prominent when the portal venous system is obstructed inside the liver.

The venous drainage from the liver is into the *right and left hepatic veins* which emerge from the back of the liver and at once enter the inferior vena cava very near its point of entry into the right atrium.

Lymphatic vessels terminate in small groups of glands around the porta hepatis. Efferent vessels drain into glands around the coeliac axis. Some superficial hepatic lymphatics pass through the diaphragm in the falciform ligament and finally reach the mediastinal glands. Another group accompanies the inferior vena cava into the thorax and ends in a few small glands around the intrathoracic portion of the inferior vena cava.

The *inferior vena cava* makes a deep groove to the right of the caudate lobe about 2 cm from the midline.

The *gallbladder* lies in a fossa extending from the inferior border of the liver to the right end of the porta hepatis.

The liver is completely covered with peritoneum, except in three places. It comes into direct contact with the diaphragm through the bare area which lies to the

right of the fossa for the inferior vena cava. The other areas without peritoneal covering are the fossae for the inferior vena cava and gallbladder.

The liver is kept in position by peritoneal ligaments and by the intra-abdominal pressure transmitted by the tone of the muscles of the abdominal wall.

Functional liver anatomy: sectors and segments

Based on the external appearances described earlier, the liver has a right and left lobe separated along the line of insertion of the falciform ligament. This separation, however, does not correlate with blood supply or biliary drainage. A *functional anatomy* is now recognized based upon vascular and biliary anatomy. The *Couinaud* classification [5] defines eight segments (segments I–IV in the left lobe, V–VIII in the right lobe), while the *Bismuth* classification [6] divides the liver into four sectors. These can be correlated with results seen with imaging techniques.

The main portal vein divides into right and left branches and each of these supplies two further subunits (variously called sectors). The sectors on the right side are anterior and posterior and, in the left lobe, medial and lateral.




Fig. 1.4 The sectors of the human liver.

and lateral – giving a total of four sectors (Fig. 1.4). Using this definition, the right and left side of the liver are divided not along the line of the falciform ligament, but along a slightly oblique line to the right of this, drawn from the inferior vena cava above to the gallbladder bed below. The right and left side are independent with regard to portal and arterial blood supply, and bile drainage. Three planes separate the four sectors and contain the three major hepatic vein branches.

Closer analysis of these four hepatic sectors produces a further subdivision into segments (Fig. 1.5). The right anterior sector contains segments V and VIII; right posterior sector, VI and VII; left medial sector, IV; left lateral sector, II and III. There is no vascular anastomosis between the macroscopic vessels of the segments but communications exist at the sinusoidal level. Segment I, the equivalent of the caudate lobe, is separate from the other segments and does not derive blood directly from the major portal branches or drain by any of the three major hepatic veins.

This functional anatomical classification allows interpretation of radiological data and is of importance to the surgeon planning a liver resection. There are wide variations in portal and hepatic vessel anatomy which can be demonstrated by spiral computed tomography (CT) and magnetic resonance imaging (MRI) reconstruction [7].

Anatomical abnormalities of the liver

These are being increasingly diagnosed with more widespread use of CT and ultrasound scanning.

Accessory lobes. The livers of the pig, dog, and camel are divided into distinct and separate lobes by strands of connective tissue. Occasionally, the human liver may




Fig. 1.5 Schematic representation of the functional anatomy of the liver. Three main hepatic veins (dark blue) divide the liver into four sectors, each of them receiving a portal pedicle; hepatic veins and portal veins are intertwined as the fingers of two hands [6].

show this reversion and up to 16 lobes have been reported. This abnormality is rare and without clinical significance. The lobes are small and usually on the undersurface of the liver so that they are not detected clinically but are noted incidentally at scanning, operation, or necropsy. Rarely, they are intrathoracic [8]. An accessory lobe may have its own mesentery containing hepatic artery, portal vein, bile duct, and hepatic vein. This may twist and demand surgical intervention.

Ectopic liver. Small nodules of normal liver derived from the embryologic hepatic bud may be found in less than 1% of laparoscopies and autopsies near the gallbladder, hepatic ligaments, gastrosplenic ligament, omentum, retroperitoneum, and thorax. These may give rise to hepatocellular carcinoma [9,10].

Riedel's lobe. This is fairly common and is a downward tongue-like projection of the right lobe of the liver [11]. It is a simple anatomical variation; it is not a true accessory lobe. The condition is more frequent in women. It is detected as a mobile tumour on the right side of the abdomen which descends with the diaphragm on inspiration. It may come down as low as the right iliac region. It is easily mistaken for other tumours in this area, especially a viscerotopic right kidney. It does not cause symptoms and treatment is not required. Rarely, it is a site for metastasis or primary hepatocellular carcinoma. Scanning may be used to identify Riedel's lobe and other anatomical abnormalities.

Cough furrows on the liver. These are vertical grooves on the convexity of the right lobe. They are one to six in number and run anteroposteriorly, being deeper posteriorly. These represent diaphragmatic sulci and fissures produced by pressure exerted by diaphragmatic muscle on peripheral structurally weak liver parenchymal zones associated with watershed vascular distribution [12]. Chronic cough produces such pressure.

Corset liver. This is a horizontal fibrotic furrow or pedicle on the anterior surface of one or both lobes of the liver just below the costal margin [13]. The mechanism is unknown, but it affects elderly women who have worn corsets for many years. It presents as an abdominal mass in front of and below the liver and is isodense with the liver. It may be confused with a hepatic tumour.

Lobar atrophy. Interference with the portal supply or biliary drainage of a lobe may cause atrophy. There is usually hypertrophy of the opposite lobe. Left lobe atrophy found at post-mortem or during scanning is not uncommon and is probably related to reduced blood supply via the left branch of the portal vein. The lobe is decreased in size with thickening of the capsule, fibrosis, and prominent biliary and vascular markings. The vascular problem may date from the time of birth. Loss of left lobe parenchyma in this instance develops by the process of ischaemic extinction due to impaired flow from the affected large portal vein branch. Replacement

fibrosis ensues. This large vessel extinction process should be distinguished from cirrhosis in which the entire liver is affected by numerous intrahepatic and discrete extinction lesions, which affect small hepatic veins and portal vein branches during the course of inflammation and fibrosis. Hence, in cirrhosis the entire liver surface is diffusely converted to regenerative parenchymal nodules surrounded by fibrosis.

Obstruction to the right or left hepatic bile duct by benign stricture or cholangiocarcinoma is now the most common cause of lobar atrophy [14]. The alkaline phosphatase is usually elevated. The bile duct may not be dilated within the atrophied lobe. Relief of obstruction may reverse the changes if cirrhosis has not developed. Distinction between a biliary and portal venous aetiology may be made using technetium-labelled iminodiacetic acid (IDA) and colloid scintiscans. A small lobe with normal uptake of IDA and colloid is compatible with a portal aetiology. Reduced or absent uptake of both isotopes favours biliary disease.

Agensis of the right lobe [15]. This rare lesion may be an incidental finding associated, probably coincidentally, with biliary tract disease and also with other congenital abnormalities. It can cause presinusoidal portal hypertension. The other liver segments undergo compensatory hypertrophy. It must be distinguished from lobar atrophy due to cirrhosis or hilar cholangiocarcinoma.

Situs inversus (SI). In the exceedingly rare *SI totalis* or *abdominalis* the liver is located in the left hypochondrium and may be associated with other anomalies including biliary atresia, polysplenia syndrome, aberrant hepatic artery anatomy, and absent portal vein. Hepatic surgery (partial hepatectomy, liver transplantation) is feasible, but complex. Other conditions associated with displacement of the liver from its location in the right upper quadrant include *congenital diaphragmatic hernias*, *diaphragmatic eventration*, and *omphalocele*.

Anatomical abnormalities of the gallbladder and biliary tract are discussed in Chapter 14.

Anatomy of the biliary tract (Fig. 1.6)

The *right* and *left hepatic ducts* emerge from the liver and unite in the porta hepatis to form the *common hepatic duct*. This is soon joined by the *cystic duct* from the gallbladder to form the common bile duct.

The *common bile duct* runs between the layers of the lesser omentum, lying anterior to the portal vein and to the right of the hepatic artery. Passing behind the first part of the duodenum in a groove on the back of the head of the pancreas, it enters the second part of the duodenum. The duct runs obliquely through the posteromedial wall, usually joining the main pancreatic duct to form the *ampulla of Vater* (c. 1720). The ampulla makes the




Fig. 1.6 Gallbladder and biliary tract.

mucous membrane bulge inwards to form an eminence, the *duodenal papilla*. In about 10–15% of subjects the bile and pancreatic ducts open separately into the duodenum.

The dimensions of the common bile duct depend on the technique used. At operation it is about 0.5–1.5 cm in diameter. Using ultrasound the values are less, the common bile duct being 2–7 mm, with values greater than 7 mm being regarded as abnormal. Using endoscopic cholangiography, the duct diameter is usually less than 11 mm, although after cholecystectomy it may be more in the absence of obstruction.

The duodenal portion of the common bile duct is surrounded by a thickening of both longitudinal and circular muscle fibres derived from the intestine. This is called the *sphincter of Oddi* (c. 1887).

The *gallbladder* is a pear-shaped bag 9 cm long with a capacity of about 50 mL. It always lies above the transverse colon, and is usually next to the duodenal cap overlying, but well anterior to, the right renal shadow. The fundus is the wider end and is directed anteriorly; this is the part palpated when the abdomen is examined. The body extends into a narrow neck which continues into the cystic duct. The *valves of Heister* are spiral folds of mucous membrane in the wall of the cystic duct and neck of the gallbladder. *Hartmann's pouch* is a sacculatation at the neck of the gallbladder; this is a common site for a gallstone to lodge.

The mucosa is in delicate, closely woven folds; instead of glands there are indentations of mucosa which usually lie superficial to the muscle layer. Increased intraluminal pressure in chronic cholecystitis results in formation of branched, diverticula-like invaginations of the mucosa which reach into the muscular layer, termed *Rokitansky–Aschoff sinuses*. There is no submucosa or muscularis mucosae. The gallbladder wall consists of a loose connective tissue lamina propria and muscular layer containing circular, longitudinal, and oblique muscle bundles without definite layers, the muscle being particularly well developed in the neck and fundus. The outer layers are the subserosa and serosa. The distensible normal gallbladder fills with bile and bile acids secreted by the liver, concentrates the bile through absorption of water and electrolytes and with meals contracts under the influence of cholecystikinin (acting through preganglionic cholinergic nerves) to empty bile into the duodenum.

Blood supply. The gallbladder receives blood from the *cystic artery*. This branch of the hepatic artery is large, tortuous, and variable in its anatomical relationships. Smaller blood vessels enter from the liver through the gallbladder fossa. The venous drainage is into the *cystic vein* and thence into the portal venous system. Attention to the vascular-biliary anatomy in the reference area known as *Calot's triangle* (bordered by the cystic duct, common hepatic duct, and lower edge of the liver) reduces the risk of vascular injuries and potential biliary strictures. Most bile duct injuries occur at cholecystectomy (incidence of <1.3% for either open or laparoscopic cholecystectomy). After liver transplantation 10–33% of patients may develop biliary complications, of which bile duct stricture is the most important.

The arterial blood supply to the supraduodenal bile duct is generally by two main (axial) vessels which run beside the bile duct. These are supplied predominantly by the retroduodenal artery from below, and the right hepatic artery from above, although many other vessels contribute. This pattern of arterial supply would explain why vascular damage results in bile duct stricturing [16].

Lymphatics. There are many lymphatic vessels in the submucous and subperitoneal layers. These drain through the cystic gland at the neck of the gallbladder to glands along the common bile duct, where they anastomose with lymphatics from the head of the pancreas.

Nerve supply. The gallbladder and bile ducts are liberally supplied with nerves, from both the parasympathetic and the sympathetic system.

Surface marking (Fig. 1.7, Fig. 1.8)

Liver. The upper border of the right lobe is on a level with the 5th rib at a point 2 cm medial to the right midclavicular line (1 cm below the right nipple). The upper border

of the left lobe corresponds to the upper border of the 6th rib at a point in the left midclavicular line (2 cm below the left nipple). Here only the diaphragm separates the liver from the apex of the heart.

The lower border passes obliquely upwards from the 9th right to the 8th left costal cartilage. In the right nipple line it lies between a point just under to 2 cm below the costal margin. It crosses the midline about midway between the base of the xiphoid and the umbilicus and the left lobe extends only 5 cm to the left of the sternum.

Gallbladder. Usually the fundus lies at the outer border of the right rectus abdominis muscle at its junction with the right costal margin (9th costal cartilage) (Fig. 1.8). In an obese subject it may be difficult to identify the outer border of the rectus sheath and the gallbladder may then be located by the Grey–Turner method. A line is drawn from the left anterior superior iliac spine through the umbilicus; its intersection with




Fig. 1.7 The surface marking of the liver.

the right costal margin indicates the position of the gallbladder. These guidelines depend upon the individual's build. The fundus may occasionally be found below the iliac crest.

Methods of examination

Liver. The lower edge should be determined by palpation just lateral to the right rectus muscle. This avoids mistaking the upper intersection of the rectus sheath for the liver edge.

The liver edge moves 1–3 cm downwards with deep inspiration. It is usually palpable in normal subjects inspiring deeply. The edge may be tender, regular or irregular, firm or soft, thickened or sharp. The lower edge may be displaced downwards by a low diaphragm, for instance in emphysema. Movements may be particularly great in athletes or singers. Some patients with practice become very efficient at 'pushing down' the liver. The normal spleen can become palpable in similar fashion. Common causes of a liver palpable below the umbilicus are malignant deposits, polycystic or Hodgkin disease, amyloidosis, congestive cardiac failure, and gross fatty change. Rapid change in liver size may occur when congestive cardiac failure is corrected, cholestatic jaundice relieved, or when severe diabetes is controlled. The surface can be palpated in the epigastrium and any irregularity or tenderness noted. An enlarged caudate lobe, as in the Budd–Chiari syndrome or with some cases of cirrhosis, may be palpated as an epigastric mass.

Pulsation of the liver, usually associated with tricuspid valvular incompetence, is felt by manual palpation with one hand behind the right lower ribs posteriorly and the other anteriorly on the abdominal wall.

The upper edge is determined by fairly heavy percussion passing downwards from the nipple line. The lower edge is recognized by very light percussion passing upwards from the umbilicus towards the costal margin. Percussion is a




Fig. 1.8 Surface markings of the gallbladder. Method I: the gallbladder is found where the outer border of the right rectus abdominis muscle intersects the 9th costal cartilage. Method II: a line drawn from the left anterior superior iliac spine through the umbilicus intersects the costal margin at the site of the gallbladder.

valuable method of determining liver size and is the only clinical method of determining a small liver.

The anterior liver span is obtained by measuring the vertical distance between the uppermost and lowermost points of hepatic dullness by percussion in the right midclavicular line. This is usually 12–15 cm. Direct percussion is as accurate as ultrasound in estimating liver span [17].

Friction may be palpable and audible, usually due to recent biopsy, tumour, or perihepatitis. The venous hum of portal hypertension is audible between the umbilicus and the xiphisternum. An arterial murmur over the liver may indicate a primary liver cancer or acute alcoholic hepatitis.

Gallbladder. The gallbladder is palpable only when it is distended. It is felt as a pear-shaped cystic mass usually about 7 cm long. In a thin person, the swelling can sometimes be seen through the anterior abdominal wall. It moves downwards on inspiration and is mobile laterally but not downwards. The swelling is dull to percussion and directly impinges on the parietal peritoneum, so that the colon is rarely in front of it. Gallbladder dullness is continuous with that of the liver.

Abdominal tenderness should be noted. Inflammation of the gallbladder causes a positive *Murphy's sign*. This is the inability to take a deep breath when the examining fingers are hooked up below the liver edge. The inflamed gallbladder is then driven against the fingers and the pain causes the patient to catch their breath.

The enlarged gallbladder must be distinguished from a *viscerototic right kidney*. This, however, is more mobile, can be displaced towards the pelvis and has the resonant colon anteriorly. A *regenerative* or *malignant nodule* feels much firmer.

Imaging. A plain film of the abdomen, including the diaphragms, may be used to assess liver size and in particular to decide whether a palpable liver is due to actual enlargement or to downward displacement. On moderate inspiration the normal level of the diaphragm, on the right side, is opposite the 11th rib posteriorly and the 6th rib anteriorly.

Ultrasound, CT, or MRI can be used to study liver size, shape, and content.

Microanatomy of the liver

For over a century, many models of liver substructure have been proposed [18]. The most popular of these is the *lobule* introduced by Kiernan in 1833 as the basic architectural unit, based on pig dissections [19]. He described circumscribed, hexagonal lobules consisting of a central tributary of the hepatic vein (central vein) and at the periphery a portal tract containing the bile

duct, portal vein radicle, and hepatic artery branch. Cords (plates) of liver cells and blood-containing sinusoids extend between these two systems. The lobule has foundations in pig, camel, raccoon, and polar bear livers, in which such hexagonal units are surrounded by interlobular connective tissue septa [20]. Such septa have no counterparts in human liver.

Stereoscopic reconstructions and scanning electron microscopy have shown the human liver as cords of liver cells radiating from a central vein, and interlaced in orderly fashion by sinusoids (Figs 1.9, 1.10). The terminal branches of the portal vein discharge their blood into the sinusoids and the direction of flow is determined by the higher pressure in the portal vein than in the central vein (or terminal hepatic venule) – see later.

The *portal tracts* are small connective tissue islands containing triads composed of the portal vein radicle, the hepatic arteriole, and bile duct (Fig. 1.11). Portal tracts are surrounded by a limiting plate of liver cells. Histological sections of normal liver show portal tracts containing dyads as frequently as triads, with the portal vein being the most frequently absent element. Within each linear centimetre of liver tissue obtained at biopsy there are usually two interlobular bile ducts, two hepatic arteries and one portal vein per portal tract, with six full portal triads [21].

The liver has to be divided *functionally*. Traditionally, the unit is based on a central hepatic vein and its surrounding liver cells. However, Rappaport [22] envisages a series of functional *acini*, each centred on the portal tract with its terminal branch of portal vein, hepatic artery, and bile duct (zone 1) (Fig. 1.12, Fig. 1.13). These interdigitate, mainly perpendicularly, with terminal hepatic veins of adjacent acini. The circulatory peripheries of acini (adjacent to terminal hepatic veins) (zone 3) suffer most from injury, whether viral, toxic, or anoxic. Bridging necrosis may extend from the periphery (acinar zone 1) to zone 3. The regions closer to the axis formed by afferent vessels and bile ducts survive longer and may later form the core from which regeneration will proceed. The contribution of each acinar zone to liver cell regeneration depends on the acinar location of damage [22].

The liver cells (*hepatocytes*) comprise about 60% of the liver. They are polygonal and approximately 30 µm in diameter. The nucleus is single or, less often, multiple and divides by mitosis. The lifespan of liver cells is about 150 days in experimental animals. The hepatocyte has three surfaces: one facing the sinusoid and space of Disse, the second facing the canaliculus, and the third facing neighbouring hepatocytes (Fig. 1.14). There is no basement membrane.

The sinusoids are lined by endothelial cells with small pores (fenestrae) for macromolecule diffusion from




Fig. 1.9 The structure of the normal human liver.




Fig. 1.10 Normal hepatic histology. H, terminal hepatic vein; P, portal tract. (H & E, $\times 60$.)




Fig. 1.11 Normal portal tract. A, hepatic artery; B, bile duct; P, portal vein. (H & E.)

blood to hepatocytes. On the vascular side of the sinusoids are the phagocytic cells of the reticulo-endothelial system (Kupffer cells) and pit cells (NK or natural killer cells) which are cytotoxic lymphoid cells involved in surveillance for tumour cells and viral infections [23,24].

There are approximately 202×10^3 cells in each milligram of normal human liver, of which 171×10^3 are parenchymal and 31×10^3 littoral (sinusoidal, including Kupffer cells).

The *space of Disse* between hepatocytes and sinusoidal endothelial cells contains a few collagen fibrils and the hepatic stellate cells, which have also been called fat-storing cells, Ito cells, and lipocytes. These cells store vitamin A and when activated in disease become collagen-synthesizing myofibroblasts. The *hepatic lymphatics* are found in the periportal connective tissue and are lined throughout by endothelium. Tissue fluid seeps through the endothelium into the lymph vessels.