

Chioma M. Okeoma *Editor*

Chikungunya Virus

Advances in Biology, Pathogenesis, and
Treatment

 Springer

Chikungunya Virus

Chioma M. Okeoma
Editor

Chikungunya Virus

Advances in Biology, Pathogenesis,
and Treatment

 Springer

Editor

Chioma M. Okeoma
Department of Microbiology
University of Iowa
Iowa City, IA, USA

Interdisciplinary Program in Molecular
and Cellular Biology
University of Iowa
Iowa City, IA, USA

ISBN 978-3-319-42956-4 ISBN 978-3-319-42958-8 (eBook)

DOI 10.1007/978-3-319-42958-8

Library of Congress Control Number: 2016960576

© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

This Springer imprint is published by Springer Nature
The registered company is Springer International Publishing AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Preface

Chikungunya virus (CHIKV) is a mosquito-transmitted virus associated with increased morbidity and causes a debilitating chronic musculoskeletal disease in infected humans. The mosquito genus *Aedes* is responsible for CHIKV transmission. *Aedes* mosquitoes were previously restricted to the tropical and subtropical countries of the world, but increased globalization has resulted in worldwide spread of CHIKV-transmitting *Aedes* mosquitoes, including the species *Aedes aegypti* and *Aedes albopictus*. Despite precautions being taken to contain the spread of CHIKV, the number of cases of traveler-associated and locally transmitted CHIKV keep increasing in many countries. In some countries, including the United States, CHIKV infection is a nationally notifiable condition reportable to government health protection agencies, such as the US Centers for Disease Control and Prevention. Sadly, there are no vaccines or effective therapies for CHIKV infection, leaving infected people to rely on their immune systems to fight the disease. Since the re-emergence of CHIKV in 1941, exciting discoveries have been made in various aspects of CHIKV research. We have done our best to report the current state of knowledge of CHIKV; however, much work remains to be done to understand fully the fundamentals of CHIKV interaction with the environment, the vector, and the hosts.

Chikungunya Virus is the first book of its kind and consists of 12 chapters written by leading experts in the broad areas of CHIKV epidemiology, CHIKV biology, mechanisms of infection and pathogenesis, host response to infection, and clinical syndromes. These chapters are independent but interrelated. The chapter on clinical syndromes highlights the complexity of CHIKV infection in patients and the current approach to managing CHIKV disease in the absence of definitive therapy. Therefore, *Chikungunya Virus* will be of great interest to a wide audience and is intended for researchers, educators, postdoctoral and medical fellows, graduate and undergraduate students, health practitioners, and other public health officials.

Iowa City, IA, USA

Chioma M. Okeoma

Contents

Clinical Syndrome and Therapy	1
Sylvie Abel and André Cabié	
Host Response and Mechanisms of Subversion of Chikungunya	19
Lisa F.P. Ng and Pierre Roques	
Epidemiological History of Chikungunya Virus	33
Ann M. Powers	
Vaccines Against Chikungunya Virus Infection	45
Karl Ljungberg, Beate M. Kümmerer, Pierre Roques, Mariano Esteban, Andres Merits, and Peter Liljeström	
Function of Chikungunya Virus Structural Proteins	63
Stefan W. Metz and Gorben P. Pijlman	
Functions of Chikungunya Virus Nonstructural Proteins	75
Tero Ahola and Andres Merits	
Interaction of Chikungunya Virus with the Mosquito Vector	99
Konstantin Tsetsarkin, Lark L. Coffey, and Scott C. Weaver	
Chikungunya Virus Entry and Replication	127
Nathalie Chazal and Laurence Briant	
Chikungunya Virus-Induced Autophagy and Apoptosis	149
Delphine Judith, Thérèse Couderc, and Marc Lecuit	
Chikungunya Virus Pathogenesis	161
David W. Hawman and Thomas E. Morrison	
Restriction Factors and Chikungunya Virus	185
Wadie D. Mahauad-Fernandez and Chioma M. Okeoma	

Contributors

Sylvie Abel Service de Maladies Infectieuses et Tropicales, Centre Hospitalier Universitaire de Martinique, Martinique, France

Tero Ahola Department of Food and Environmental Sciences, University of Helsinki, Helsinki, Finland

Laurence Briant CPBS, Centre d'études d'agents Pathogènes et Biotechnologies pour la Santé, UMR5236 CNRS, Université Montpellier, Montpellier, France

André Cabié Service de Maladies infectieuses et tropicales, Centre Hospitalier Universitaire de Martinique, Martinique, France

Nathalie Chazal CPBS, Centre d'études d'agents Pathogènes et Biotechnologies pour la Santé, UMR5236 CNRS, Université Montpellier, Montpellier, France

Lark L. Coffey Center for Vectorborne Diseases and Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine, University of California, Davis, CA, USA

Thérèse Couderc Institut Pasteur, Biology of Infection Unit, Paris, France
Inserm U1117, Paris, France

Mariano Esteban Department of Molecular and Cellular Biology, Centro Nacional de Biotecnología, CSIC, Madrid, Spain

David W. Hawman Department of Immunology and Microbiology, University of Colorado School of Medicine, Aurora, CO, USA

Delphine Judith Molecular Cell Biology of Autophagy Group, Francis Crick Institute, London, UK

Institut Pasteur, Biology of Infection Unit, Paris, France
Inserm U1117, Paris, France

Beate M. Kümmerer University of Bonn Medical Centre, Bonn, Germany

Marc Lecuit Department of Infectious Diseases and Tropical Medicine, Paris Descartes University, Necker-Enfants Malades University Hospital, Paris, France
Institut Pasteur, Biology of Infection Unit, Paris, France
Inserm U1117, Paris, France

Peter Liljeström Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

Karl Ljungberg Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

Wadie D. Mahauad-Fernandez Department of Microbiology, University of Iowa, Iowa City, IA, USA
Interdisciplinary Program in Molecular and Cellular Biology, University of Iowa, Iowa City, IA, USA

Andres Merits Institute of Technology, University of Tartu, Tartu, Estonia

Stefan W. Metz Department of Microbiology and Immunology, University of North Carolina, School of Medicine, Chapel Hill, NC, USA
Laboratory of Virology, Wageningen University, Wageningen, The Netherlands

Thomas E. Morrison Department of Immunology and Microbiology, University of Colorado School of Medicine, Aurora, CO, USA

Lisa F.P. Ng Singapore Immunology Network, A*STAR, Singapore, Singapore

Chioma M. Okeoma Department of Microbiology, University of Iowa, Iowa City, IA, USA
Interdisciplinary Program in Molecular and Cellular Biology, University of Iowa, Iowa City, IA, USA

Gorben P. Pijlman Laboratory of Virology, Wageningen University, Wageningen, The Netherlands

Ann M. Powers Division of Vector Borne Diseases, Centers for Disease Control and Prevention, Fort Collins, CO, USA

Pierre Roques Immuno-Virology Department, Institute of Emerging Disease and Innovative Therapy, CEA, Fontenay aux Roses, France
University Paris-Sud XI, UMR-E1, Orsay, France

Konstantin Tsetsarkin National Institute of Allergy and Infectious Diseases, Laboratory of Infectious Diseases, Neurotropic Flaviviruses Section, Bethesda, MD, USA

Scott C. Weaver Institute for Human Infections and Immunity and Center for Tropical Diseases, University of Texas Medical Branch, Galveston, TX, USA

Clinical Syndrome and Therapy

Sylvie Abel and André Cabié

Introduction

Chikungunya virus (CHIKV), “that which bends up” in the Makonde dialect is an emerging alphavirus transmitted to humans by *Aedes* mosquitoes (*Ae. aegypti*, *Ae. albopictus*). CHIKV emergence in all the tropical zones at the beginning of the twenty-first century allowed experts to discover the polymorphism of its clinical manifestations (Weaver and Lecuit 2015).

The female mosquito becomes infected after feeding on blood of a viremic person (viremia for 5–7 days after onset of clinical signs). The virus replicates in the mosquito for a few days, and then the mosquito can transmit the virus to another person, throughout its life (Fig. 1; Schwartz and Albert 2010). Mother-to-child transmission can occur during childbirth when the mother is viremic (Gérardin et al. 2008). Although direct person-to-person transmission has not been reported, nosocomial transmission most probably occurs following blood transfusion or needlestick injury (Gallian et al. 2014; Parola et al. 2006).

CHIKV infection is most often symptomatic ($\approx 80\%$ of cases); the symptoms last from a few days to several years depending on the case. In France, experts have defined three clinical stages (Simon et al. 2015): acute stage (from the first day on which the first symptoms appear (D1) up to day 21 (D21)); post-acute stage (from D21 to the end of the third month); and chronic stage (after 3 months; Fig. 1). This time staging takes into account the pathogenic, clinical, and therapeutic variations over time. The post-acute stage and a fortiori the chronic stage are not observed in all patients. The mortality rate of CHIKV is comparable to that of seasonal influenza (≈ 0.01 to 0.1%), and is mainly related to the patient’s age (increased over 75 years) and/or to chronic comorbidities (Schwartz and Albert 2010).

S. Abel • A. Cabié (✉)

Service de Maladies infectieuses et tropicales, Centre Hospitalier
Universitaire de Martinique, Martinique, France
e-mail: andre.cabie@chu-fortdefrance.fr; andrecabie@orange.fr




Fig. 1 Transmission cycle and natural history of chikungunya. D1 defined as the day of symptom onset

Acute Stage

Clinical Manifestations and Symptoms

Arboviral Syndrome

In contrast to other arboviral diseases, CHIKV infection is symptomatic in more than 70 % of cases, and ranges from 72 to 96 % (Appassakij et al. 2013). In symptomatic patients the mean incubation period is 3 days, range 1–12 days (Burt et al. 2012). In the common presentation, chikungunya is a rapid-onset febrile disease with no prodromal phase (Thiberville et al. 2013). High-grade fever occurs suddenly, along with inflammatory arthralgia and arthritis with periarticular edema and sometimes severe pain (Fig. 2). Joint pain is mostly polyarticular, bilateral, symmetrical, and occurs mainly in peripheral joints (wrists, ankles, and phalanges) and some large joints (shoulders, elbows, and knees). Sometimes arthralgia appears a few hours before the onset of fever. Other typical symptoms are: myalgia, headache, backache, macular to maculopapular rash, frequently associated with cutaneous pruritus (palms and soles) and edema of the face, lymphadenopathy. The rash appears after fever onset and is typically maculopapular involving the trunk and extremities but can also involve palms, soles, and the face (Fig. 3). External ear redness has been observed, and this may reflect chondritis and is evocative of CHIKV infection (Fig. 4) (Javelle et al. 2014). Fever and cutaneous

Fig. 2 Acute chikungunya: arthritis and distal edema



Fig. 3 Acute chikungunya: maculopapular rash and palm erythema



Fig. 4 Acute chikungunya: rash of the face and external ear redness




Table 1 Comparison of clinical and biological manifestations of dengue, chikungunya, and Zika virus infection

	Chikungunya	Dengue	Zika virus infection
Fever	+++	+++	++
Myalgia	+	+++	++
Arthralgia	+++	++	++
Retro-orbital pain	+	+++	+
Skin rash	+++	++	+++
Nonpurulent conjunctivitis	+	0	+++
Arthritis/edema	+++	0	++
Hypotension	+	+++	+
Minor bleeding	±	++	±
Lymphadenopathy	++	++	+
Thrombocytopenia	+	+++	±
Lymphopenia	+++	+++	±

rash last 3–5 days (viremic period), but articular manifestations may last 2–3 weeks in some patients (Thiberville et al. 2013). Asthenia and anorexia are common after regression of fever. The main laboratory finding is lymphopenia, which is closely associated with viremia when the lymphocyte count is less than 1000 per cubic millimeter. Other laboratory abnormalities include moderate thrombocytopenia, increased levels of aspartate aminotransferase, alanine aminotransferase, and creatine phosphokinase, and mild increase in C-reactive protein levels (about 50–60 mg/L). The clinical presentation is similar to that of other arboviruses (dengue, Zika virus infection) and may pose diagnostic difficulties (Table 1; Ios et al. 2014; Simon et al. 2011; Staples et al. 2009).

Atypical Manifestations

Atypical presentations are observed in 0.5 % of cases, mainly in vulnerable patients (young children, elderly patients, chronic alcohol abusers, patients presenting with chronic medical conditions including systemic lupus; Economopoulou et al. 2009; Rajapakse et al. 2010). Atypical presentations include hyperalgetic symptoms, gastrointestinal symptoms (diarrhea, vomiting, abdominal pain), and neurological symptoms (confusion, optic neuritis), damage to mucous membranes (oral or genital ulceration, conjunctivitis), and malaise (hypotension, dysautonomia). CHIKV can directly induce severe atypical presentations (rhabdomyolysis, bullous dermatosis, fulminant hepatitis, encephalitis or encephalopathy, Guillain-Barré syndrome, polyneuropathy, myocarditis, multiple organ failure; Betancur et al. 2016; Das et al. 2010; Farnon et al. 2008; Lemant et al. 2008; Simon et al. 2008). More frequently, it causes decompensation of chronic cardiac, respiratory, renal, systemic (lupus), and metabolic (diabetes) diseases, or various complications (dehydration, thromboembolism, loss of autonomy). The risk of drug toxicity by overdose

(self-medication) or drug interaction is high for acetaminophen as well as for other analgesics, anti-inflammatory drugs, long-term treatments, and traditional medicines used for self-medication.

Neonatal Chikungunya

CHIKV infection has not been linked to increased risk of miscarriage, fetal death in utero, or birth defects. However maternal–neonatal transmission can occur in viremic women during childbirth. Fifty percent of neonates are infected when they are born the day before or 5 days after the mother’s first symptoms (Ramful et al. 2007). Neonatal chikungunya can either be congenital or neonatal (by mosquito bite after birth). Infected neonates exhibit atypical clinical presentation (fever, difficulty to breast-feed, and pain) occurring after a median incubation period of 4 days (3–7 days; Gérardin et al. 2008; Ramful et al. 2007). The main laboratory findings are thrombocytopenia, lymphopenia, and moderate increased levels of aspartate aminotransferase and alanine aminotransferase in blood. Severe manifestations occur in 50% of cases, which include encephalopathy with progressive cerebral edema, hemodynamic disorders inducing severe sepsis, hemorrhagic complications due to intravascular coagulation, and cardiomyopathy. The mortality rate of severe presentations is 50% and the risk of post-encephalopathy psychomotor sequelae is important (Gérardin et al. 2014).

The disease presentation in infants and children is often similar to that of adults. Nevertheless, some atypical or complicated presentations have been reported including hyperalgesia resistant to analgesic treatment, extensive bullous rash, hemodynamic disorders, dehydration, food intolerance, seizures, and meningeal syndrome.

Clinical Assessment

The first step is to discuss CHIKV infection in a patient with an acute onset presentation. A suspected case of acute chikungunya is defined by the combination of fever >38.5 °C and sudden onset of debilitating joint pain without any infectious focus. However, less symptomatic or atypical presentations have been reported. Physicians should be aware of other possible diagnoses in tropical areas: dengue, Zika virus infections, meningitis, malaria, and leptospirosis. The absence of joint involvement, predominance of myalgia, a hypotensive or bleeding trend, abdominal pain, and fever for more than 5 days may justify search for other diagnoses.

The clinical step allows identifying proven clinical signs of severity, atypical and/or complicated presentations (intense pain, organ failure, bleeding, dehydration, decompensation of comorbidity, thrombosis), pregnant women, patients at risk of severe presentations (neonates, children with a history of febrile convulsion, elderly patients, chronic disease treatment, social isolation), and guiding the patient triage (hospital admission or consultation, outpatient management; Fig. 5).




Fig. 5 Acute chikungunya: management of a suspected acute chikungunya

Some pediatric presentations may be atypical or complicated (hyperalgesia despite analgesic treatment, extensive bullous rash, hemodynamic disorders, dehydration, food intolerance, seizures, and meningeal syndrome) and should be referred to the emergency unit (Economopoulou et al. 2009; Ramful et al. 2007).

The suspicion of CHIKV infection in pregnant patients requires screening for signs of severity including fever >39 °C, neurological disorders, bleeding, uterine contractions, inability to eat, poor global health status, and/or alteration of the fetal heart rate pattern (FHRP) after 28 weeks. Any sign of severity requires emergency hospitalization, at best in obstetrics and neonatal resuscitation. A hospital consultation is recommended for any case of suspected chikungunya in the last 3 months of pregnancy. CHIKV infection suspected on the basis of a common presentation should be confirmed by ruling out other causes of potentially severe fever, according to the

clinical presentation, e.g., listeria, pyelonephritis, toxoplasmosis, rubella, malaria, dengue, Zika virus infection, and by recording the FHRP in the case of contractions, in order to define an obstetrical strategy. Furthermore, during epidemics, all patients in labor should be questioned about symptoms in the delivery room to identify any risk of CHIKV transmission for the unborn child.

Social isolation should also be taken into account to organize care, because of the great risk of rapid loss of autonomy among the weakest patients.

The clinical evaluation, in the acute stage, is sufficient to assess the impact of musculoskeletal lesions by identifying the site and the intensity of inflammatory manifestations. There is no indication for X-rays or ultrasound of the joints at this stage, except for another diagnosis.

Diagnosis of Acute CHIKV Infection

In contrast to dengue, routine biological tests are not essential for typical uncomplicated presentations in patients without any chronic disease or risk. The assessment of complete blood count, kidney and liver function, blood glucose, fluid and electrolyte level, and level of inflammation should be decided on a case-by-case basis. Screening for a differential diagnosis may justify implementing additional laboratory tests in the case of atypical clinical presentation, and complicated or abnormal outcome. Main differential diagnoses are dengue or Zika virus infection (possible coinfection), acute HIV infection, malaria, leptospirosis, sepsis, post-streptococcal immune reactions, and other acute viral infections.

The need for virological CHIKV infection confirmation depends on clinical manifestations and on the epidemic context. CHIKV infection confirmation is needed in cases of atypical or severe manifestations, in patients at risk of severe presentations (chronic diseases, extreme ages, pregnancy), in sporadic suspected cases, or in the first weeks of CHIKV emergence in a naïve population, as well as for other public health issues (study of strains, suspicion of a new focus, suspected post-epidemic cases). Conversely, in epidemic regions, diagnostic confirmation in the acute stage is not recommended during epidemics for typical cases without risk of severe presentation.

The confirmed diagnosis relies on virus detection through reverse transcriptase–polymerase chain reaction (RT-PCR) testing during the first week (Fig. 6; Simon et al. 2011). Specific antibodies detection is facilitated by the limited antigenic diversity of CHIKV and extensive cross-reactivity of the antibodies induced by different strains. Serum IgM is detectable from day 5 (and even earlier) to several months after the onset of illness and is also considered as a diagnostic parameter. Seroconversion can also be detected by an increase in IgG by a factor of 4 or more between the acute phase and convalescent phases. The tests are usually performed as follows: RT-PCR between Day 1 and Day 5, RT-PCR and serology between Day 5 and Day 7, serology alone after Day 7; viral culture is not routinely performed. The interpretation of the tests is based on the epidemiological context and clinical information provided by the clinician (time of onset of symptoms is mandatory).