

BIOMARKERS IN TOXICOLOGY

BIOMARKERS IN TOXICOLOGY

Edited by

RAMESH C. GUPTA, DVM, MVSc, PhD, DABT, FACT, FACN, FATS

*Professor and Head, Toxicology Department
Breathitt Veterinary Center
Murray State University
Hopkinsville, Kentucky
USA*



ELSEVIER

AMSTERDAM • BOSTON • HEIDELBERG • LONDON
NEW YORK • OXFORD • PARIS • SAN DIEGO
SAN FRANCISCO • SINGAPORE • SYDNEY • TOKYO

Academic Press is an imprint of Elsevier



Academic Press is an imprint of Elsevier
525 B Street, Suite 1800, San Diego, CA 92101-4495, USA
32 Jamestown Road, London NW1 7BY, UK
225 Wyman Street, Waltham, MA 02451, USA

Copyright © 2014 Elsevier Inc. All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the publisher.

Permissions may be sought directly from Elsevier's Science & Technology Rights Department in Oxford, UK: phone (+44) (0) 1865 843830; fax (+44) (0) 1865 853333; email: permissions@elsevier.com. Alternatively, visit the Science and Technology Books website at www.elsevierdirect.com/rights for further information.

Notice

No responsibility is assumed by the publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. Because of rapid advances in the medical sciences, in particular, independent verification of diagnoses and drug dosages should be made.

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library

Library of Congress Cataloging-in-Publication Data

A catalog record for this book is available from the Library of Congress

ISBN: 978-0-12-404630-6

For information on all Academic Press publications
visit our website at elsevierdirect.com

Typeset by MPS Limited, Chennai, India
www.adi-mps.com

Printed and bound in the United States of America

14 15 16 17 18 10 9 8 7 6 5 4 3 2 1

		Working together to grow libraries in developing countries
www.elsevier.com • www.bookaid.org		

*This book is dedicated to my daughter Rekha, wife Denise,
and parents the late Chandra and Triveni Gupta.*

Foreword

Merriam-Webster defines a biomarker as a distinctive biological or biologically derived indicator (as a metabolite) of a process, event, or condition (as, for example, aging, disease, or oil formation). The subject of the present book is biomarkers in toxicology, but it should be remembered that biomarkers include substances used in the detection of numerous diseases, including the autoimmune diseases, which are not generally thought of as toxicological in origin.

The earliest biomarkers of toxicological exposure date from a time before precise analytical techniques were available and include the Kayser-Fleischer ring, usually indicative of copper accumulation in the cornea in cases of Wilson's disease, as well as lead lines in the gums associated with lead toxicity. The cherry-red color noted as a (rather unreliable) clinical sign in carbon monoxide poisoning may also be described as a biomarker.

One of the earliest biomarkers relying on biochemical analytical techniques was cholinesterase, initially whole blood cholinesterase and later plasma pseudocholinesterase (butyrylcholinesterase) and red blood cell acetylcholinesterase. Cholinesterase measurements were introduced at defense laboratories after World War II as a screening test for excessive exposure to organophosphate compounds: a 20% depression in activity was considered a mandate for cessation of exposure of individuals to organophosphate nerve agents. Pseudocholinesterase and red blood cell acetylcholinesterase measurements now have numerous uses in worker protection, clinical diagnosis of poisoning, and human and experimental animal studies (including regulatory studies) in relation to the use of organophosphate and other anticholinesterase pesticides. Since World War II, biomarkers of toxicity have ballooned in importance and number in worker and consumer protection and in clinical toxicology and experimental

toxicology and are also widely used in regulation of chemicals in animal and, less commonly, human experimental studies. Toxicological biomarkers are also used in allied disciplines – for example epidemiology – and may be used to estimate population loads of exposure.

In toxicology, biomarkers are often divided into biomarkers of exposure, of effect, and of susceptibility and all of these are dealt with in this extremely wide-ranging book. The book has an initial introductory section, including discussion of rodent, minipig, primate, zebrafish, *Caenorhabditis elegans*, and alternative animal models for toxicological testing and biomarkers and pharmacokinetics, PBPK modeling, and toxicological biomarkers. Part II, Systems Toxicity Biomarkers, comprises chapters on biomarkers of toxicity in relation to all important organs and organ systems. Part III deals with biomarkers in relation to the toxicity of specific groups of compounds, and includes chapters on pesticides, as well as polychlorinated biphenyls (PCBs), polybrominated biphenyls (PBBs), and brominated flame retardants, polycyclic aromatic hydrocarbons (PAHs), bisphenol A, melamine, metals, chemical and biological warfare agents, mycotoxins, and poisonous plants. Part IV covers pharmaceuticals and nutraceuticals, with chapters on drug toxicity biomarkers, and toxicological testing, and biomarkers in the regulation of dietary supplements and nutrigenomic biomarkers. Part V covers biomarkers of petroleum products toxicity and chemical mixtures toxicity. Part VI is devoted to biomarkers of radiation, nanoparticles, carcinogens, and carcinogenesis, while Part VII deals with chapters on miscellaneous topics, including biomarkers of neurodegenerative diseases, such as Alzheimer's and Parkinson's. In addition, this section includes chapters on cytoskeletal disruption as a biomarker of developmental neurotoxicity, biomarkers of

mitochondrial dysfunction and toxicity, and biomarkers of oxidative and nitrosative stress. There are also chapters on citrulline as a biomarker of diseases and toxicities and pathological biomarkers in toxicology. Part VIII is devoted to applications of biomarkers in toxicology with ten extremely comprehensive chapters.

The 64-chapter book has an outstanding array of authors from the USA, Canada, Denmark, Finland,

Greece, Hungary, India, Italy, Japan, Portugal, South Africa, and Spain. Professor Gupta deserves our gratitude for assembling such a distinguished group of experts to produce a comprehensive book on this rapidly growing and very important field.

Timothy C Marrs
Edenbridge, UK

List of Contributors

Angela Aliberti

St. John's University, College of Pharmacy and Health Sciences, Department of Pharmaceutical Sciences, Toxicology Division, Queens, NY, USA

Patrick Allard

Department of Environmental Health Sciences and Institute for Society & Genetics, University of California, Los Angeles, Los Angeles, CA, USA

Arturo Anadón

Department of Toxicology and Pharmacology, Faculty of Veterinary Medicine, Universidad Complutense de Madrid, Madrid, Spain

Vellareddy Anantharam

Parkinson's Disorder Research Laboratory, Iowa Center for Advanced Neurotoxicology, Department of Biomedical Sciences, Iowa State University, Ames, IA, USA

Anthony E. Archibong

Department of Physiology, Meharry Medical College, Nashville, TN, USA

Adam D. Aulbach

MPI Research, Mattawan, MI, USA

Aryamitra Banerjee

Toxicology Research Laboratory, University of Illinois at Chicago, Chicago, IL, USA

Leah D. Banks

Department of Biochemistry & Cancer Biology, Meharry Medical College, Nashville, TN, USA

Frank A. Barile

College of Pharmacy and Health Sciences, Department of Pharmaceutical Sciences, Toxicology Division, St. John's University, Queens, NY, USA

Sudheer R. Beedanagari

Bristol Myers Squibb, New Brunswick, NJ, USA

Enrico Bergamaschi

Department of Clinical and Experimental Medicine, Occupational and Environmental Medicine Unit, University of Parma, Parma, Italy

Sneha P. Bhatia

Research Institute for Fragrance Materials, NJ, USA

Karyn Bischoff

Senior Extension Associate, Diagnostic Toxicologist, Cornell University, New York State Animal Health Diagnostic Center, Ithaca, NY, USA

Víctor Castellano

Department of Toxicology and Pharmacology, Faculty of Veterinary Medicine, Universidad Complutense de Madrid, Madrid, Spain

Daniel T. Chang

US Environmental Protection Agency, National Exposure Research Laboratory, Research Triangle Park, NC, USA

James J. Chen

Director of Biostatistics Branch, Division of Bioinformatics and Biostatistics, National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR, USA

Deborah Citrin

Investigator, Radiation Oncology Branch,
Center for Cancer Research, National Cancer Institute,
Bethesda, MD, USA

Rory Conolly

US Environmental Protection Agency, National
Health and Environmental Effects Research Laboratory,
Research Triangle Park, NC, USA

R.W. Coppock

Adjunct Professor, Department of Public Health
Sciences, School of Public Health University of Alberta,
Adjunct Professor, Environmental Health, Concordia
University College of Alberta, Edmonton, AB, Canada;
President and CEO, Robert W. Coppock, Toxicologist
and Associates Ltd, Vegreville, AB, Canada

Lucio G. Costa

Department of Environmental and Occupational Health
Sciences, University of Washington, Seattle, WA, USA

T.V. Damodaran

Department of Biology, North Carolina Central
University, Durham, NC, USA

Kellye K. Daniels

Director, Toxicology and Pathology, Southern Research
Institute, Birmingham, AL, USA

Curtis C. Dary

US Environmental Protection Agency, National
Exposure Research Laboratory, Las Vegas, NV, USA

Robin B. Doss

Murray State University, Toxicology Department,
Breathitt Veterinary Center, Hopkinsville, KY, USA

Filipe V. Duarte

Department of Life Sciences, Center for Neurosciences
and Cell Biology, University of Coimbra, Portugal

Margitta M. Dziwenka

Assistant Director of Operations, Health Sciences
Laboratory Animal Services, Faculty of Medicine and
Dentistry, University of Alberta, Edmonton, AB,
Canada; President and CEO, Toxalta Consulting Ltd.
Vegreville, AB, Canada

Stephen Edwards

US Environmental Protection Agency, National Health
and Environmental Effects Research Laboratory,
Research Triangle Park, NC, USA

Jorge Estévez

Instituto de Bioingeniería, Universidad Miguel
Hernández de Elche, Spain

Daniel S. Fabricant

Director, Division of Dietary Supplement Programs, Center
for Food Safety and Applied Nutrition, US Food and Drug
Administration, College Park, MD, USA

Anna M. Fan

Chief, Pesticide and Environmental Toxicology Branch,
Office of Environmental Health Hazard Assessment,
California Environmental Protection Agency,
Oakland, CA, USA

Ali Faqi

Senior Director of Developmental and Reproductive
Toxicology, MPI Research, Mattawan, MI, USA

Suzanne E. Fenton

National Toxicology Program (NTP) Laboratories
Branch, Division of the NTP, National Institute of
Environmental Health Sciences, National Institutes
of Health, Department of Health and Human Services,
Research Triangle Park, NC, USA

Vanessa A Fitsanakis

King University, Department of Biology, Bristol,
TN, USA

John Flaskos

Laboratory of Biochemistry and Toxicology,
Faculty of Veterinary Medicine, Aristotle University
of Thessaloniki, Thessaloniki, Greece

Swaran J.S. Flora

Division of Regulatory Toxicology, Defence Research
and Development, Gwalior, MP, India

Sue M. Ford

St. John's University, College of Pharmacy & Health
Sciences, Pharmaceutical Sciences Department,
Toxicology Program, Jamaica, NY, USA

Domniki Fragou

Laboratory of Forensic Medicine and Toxicology, School
of Medicine, Aristotle University of Thessaloniki, Greece

Shayne C. Gad

Gad Consulting Services, Cary, NC, USA

Niels-Christian Ganderup

Chief Scientific Officer, Ellegaard Göttingen Minipigs
A/S, Dalmose, Denmark

Dale R. Gardner

United States Department of Agriculture, Agricultural Research Service, Poisonous Plant Research Laboratory, Logan, UT, USA

Ronette Gehring

Department of Anatomy and Physiology, College of Veterinary Medicine, Kansas State University, Manhattan, KS, USA

Fernando Gil

Department of Legal Medicine and Toxicology, University of Granada School of Medicine, Granada, Spain

Saryu Goel

Associate Director & Project Leader, Preclinical & Competitive Intelligence, Supernus Pharmaceuticals, Inc. Rockville, MD, USA

Michael-Rock Goldsmith

US Environmental Protection Agency, National Exposure Research Laboratory, Research Triangle Park, NC, USA

Christopher M. Grulke

US Environmental Protection Agency, National Exposure Research Laboratory, Research Triangle Park, NC, USA

Mary Gulumian

Toxicology Department, National Institute for Occupational Health, Hematology and Molecular Department of Toxicology, National Institute of Occupational Health, Johannesburg, South Africa

P.K. Gupta

Patron and Founder, President, Society of Toxicology Founder, Editor-in-Chief, *Toxicology International* President, Academy of Sciences for Animal Welfare, Former Head of Div. Pharmacology & Toxicology, IVRI and Advisor to WHO, C- 44, Rajendra Nagar, Bareilly, UP, India

Ramesh C. Gupta

Professor and Head, Murray State University, Toxicology Department, Breathitt Veterinary Center, Hopkinsville, KY, USA

Sharon Gwaltney-Brant

Consultant, Veterinary Information Network, Adjunct Faculty, Department of Comparative Biosciences, College of Veterinary Medicine, University of Illinois at Urbana-Champaign, IL, USA

Alan J Hargreaves

School of Science and Technology, Nottingham Trent University, Nottingham, UK

Kelly L. Harris

Department of Biochemistry & Cancer Biology, Meharry Medical College, Nashville, TN, USA

Holly E. Hatfield

Department of Biochemistry, University of Cincinnati Medical School, Cincinnati, OH, USA

Diane E. Heck

Department of Environmental Health Science, School of Health Sciences and Practice, New York Medical College, Valhalla, NY, USA

Antonio F. Hernández

Department of Legal Medicine and Toxicology, University of Granada School of Medicine, Granada, Spain

Corey J. Hilmas

Chief, Dietary Supplements Regulation Implementation Branch, Division of Dietary Supplement Programs, Center for Food Safety and Applied Nutrition, US Food and Drug Administration, College Park, MD, USA

Evangelia Hondroulis

Nanobioengineering/Bioelectronics Laboratory, Department of Biomedical Engineering, Florida International University, Miami, FL, USA

Darryl B. Hood

Division of Environmental Health Science, College of Public Health, Department of Neuroscience, College of Medicine, The Ohio State University, Columbus, OH, USA

Kathryn Hudak

Research Fellow, Radiation Oncology Branch, Center for Cancer Research, National Cancer Institute, Bethesda, MD, USA

Pasi Huuskonen

School of Pharmacy/Toxicology, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland

Stewart B. Jacobson

Shin Nippon Biomedical Laboratories USA, Ltd, Everett, WA, USA

Huajun Jin

Parkinson's Disorder Research Laboratory, Iowa Center for Advanced Neurotoxicology, Department of Biomedical Sciences, Iowa State University, Ames, IA, USA

Laurie B. Joseph

Department of Pharmacology and Toxicology,
Rutgers University, Ernest Mario School of Pharmacy,
Piscataway, NJ, USA

Jun Kanno

Division of Cellular and Molecular Toxicology,
Biological Safety Research Center, National Institute
of Health Sciences, Tokyo, Japan

Anumantha G. Kanthasamy

Parkinson's Disorder Research Laboratory,
Iowa Center for Advanced Neurotoxicology,
Department of Biomedical Sciences, Iowa State
University, Ames, IA, USA

Arthi Kanthasamy

Parkinson's Disorder Research Laboratory,
Iowa Center for Advanced Neurotoxicology,
Department of Biomedical Sciences, Iowa State
University, Ames, IA, USA

Navinchandra M. Kaore

Department of Microbiology, People's College
of Medical Sciences & Research Center, Bhopal, MP ,
India

Shilpa N. Kaore

Department of Pharmacology, People's College
of Medical Sciences & Research Center, Bhopal,
MP, India

Bhupendra S. Kaphalia

Department of Pathology, University of Texas Medical
Branch, Galveston, TX, USA

Vesa Karttunen

School of Pharmacy/Toxicology, Faculty of
Health Sciences, University of Eastern Finland,
Kuopio, Finland

Greer E. Kaufman

Department of Bioanalysis, Southern Research Institute,
Birmingham, AL, USA

Ravneet Kaur

Aveley, Western Australia

Hong Duck Kim

Department of Environmental Health Science, School of
Health Sciences and Practice, New York Medical
College, Valhalla NY, USA

Prasada Rao S. Kodavanti

Neurotoxicology Branch, Toxicity Assessment Division,
NHEERL/ORD, US Environmental Protection Agency,
Research Triangle Park, NC, USA

Urmila P. Kodavanti

Environmental Public Health Division, National
Health and Environmental Effects Research Laboratory,
Office of Research and Development, US Environmental
Protection Agency, Research Triangle Park, NC, USA

George A. Kontadakis

Department of Ophthalmology, Faculty of Medicine,
University of Crete, Heraklion, Greece

Péter Krajcsi

SOLVO Biotechnology, Budaörs, Hungary

Gopala Krishna

Executive Director, Preclinical & Competitive
Intelligence, Supernus Pharmaceuticals, Inc., Rockville,
MD, USA

Kavya A. Krishna

Supernus Pharmaceuticals, Inc. Rockville, MD, USA

Maria Kummu

Department of Pharmacology and Toxicology,
Institute of Biomedicine, and Center for Arctic
Medicine, Thule Institute of Oulu, University of Oulu,
Oulu, Finland

George D. Kymionis

Department of Ophthalmology, Faculty of Medicine,
University of Crete, Heraklion, Greece

Michelle A. Lasher

Murray State University, Toxicology Department,
Breathitt Veterinary Center, Hopkinsville, KY, USA

Chen-zhong Li

Nanobioengineering/Bioelectronics Laboratory,
Department of Biomedical Engineering, Florida
International University, Miami, FL, USA

Wei-Jiun Lin

Assistant Professor, Department of Applied
Mathematics, Feng Chia University, Taiwan

Bommanna G. Loganathan

Department of Chemistry and Watershed Studies
Institute, Murray State University, Murray, KY, USA

Jarkko Loikkanen

School of Pharmacy/Toxicology, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland

Anna E. Lokshin

University of Pittsburgh Cancer Institute, Hillman Cancer Center, Department of Medicine, Department of Pathology, Department of Ob/Gyn, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA

Marcello Lotti

Professor of Medicine, Dipartimento di Scienze Cardiologiche, Toraciche e Vascolari, Università degli Studi Padova, Italy

Tzu-Pin Lu

Postdoctoral Fellow, Division of Bioinformatics and Biostatistics, National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR, USA

Yi Lu

Department of Pathology and Laboratory Medicine, University of Tennessee Health Science Center, Memphis, TN, USA

Chikezie Madu

Department of Pathology and Laboratory Medicine, University of Tennessee Health Science Center, Memphis, TN, USA

Rémi Magnan

SOLVO Biotechnology, Budaörs, Hungary

Brinda Mahadevan

Medical Safety & Surveillance, Abbott Laboratories, Columbus, OH, USA

Jane A. Mantey

Department of Neuroscience & Pharmacology, Meharry Medical College, Nashville, TN, USA

María Rosa Martínez-Larrañaga

Department of Toxicology and Pharmacology, Faculty of Veterinary Medicine, Universidad Complutense de Madrid, Madrid, Spain

Vincent P. Meador

Covance Laboratories, Inc. Global Pathology, Madison, WI, USA

Dejan Milatovic

Charlottesville, VA, USA

Pushpinder K. Multani

Stress Neurobiology Division, Department of Anesthesiology, Abramson Cancer Center, Children's Hospital of Philadelphia, Philadelphia, PA, USA

Päivi Myllynen

Center for Arctic Medicine, Thule Institute, University of Oulu, Oulu, Finland

Kirsi Myöhänen

ECHA, European Chemicals Agency, Helsinki, Finland

Rekek Negga

King University, Department of Biology, Bristol, TN, USA

Jairo Nelson

Nanobioengineering/Bioelectronics Laboratory, Department of Biomedical Engineering, Florida International University, Miami, FL, USA

Brian M. Nolen

University of Pittsburgh Cancer Institute, Hillman Cancer Center, Pittsburgh, PA, USA

Meliton N. Novilla

Shin Nippon Biomedical Laboratories USA, Ltd, Everett, WA, USA; Adjunct Associate Professor of Pathobiology, School of Veterinary Medicine, Purdue University, West Lafayette, IN, USA

Stephanie Padilla

Integrated Systems Toxicology Division, National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, Research Triangle Park, NC, USA

Carlos M. Palmeira

Department of Life Sciences, Center for Neurosciences and Cell Biology, University of Coimbra, Portugal

Kip E. Panter

United States Department of Agriculture, Agricultural Research Service, Poisonous Plant Research Laboratory, Logan, UT, USA

Markku Pasanen

School of Pharmacy/Toxicology, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland

Daniel J. Patrick

Director of Pathology, MPI Research, Mattawan, MI, USA

Sofia Pavanello

Research Scientist, Dipartimento di Scienze
Cardiologiche, Toraciche e Vascolari, Università degli
Studi Padova, Italy

Olavi Pelkonen

Department of Pharmacology and Toxicology,
Institute of Biomedicine, University of Oulu,
Oulu, Finland

Claudia Pellacani

Department of Neuroscience, University of Parma,
Parma, Italy

Michael A. Pellizzon

Research Diets, Inc., New Brunswick, NJ, USA

Andrew D. Penman

Vice President, Drug Development, Southern Research
Institute, Birmingham, AL, USA

Martin Phillips

US Environmental Protection Agency, National
Exposure Research Laboratory, Research Triangle Park,
NC, USA

Jason Pitt

Northwestern University, Department of Physiology,
Chicago, IL, USA

Argyro Plaka

Department of Ophthalmology, Faculty of Medicine,
University of Crete, Heraklion, Greece

Aramandla Ramesh

Department of Biochemistry & Cancer Biology, Meharry
Medical College, Nashville, TN, USA

Kausik Ray

Scientific Review Officer, Scientific Review Branch,
Division of Extramural Activities, National
Institute on Deafness and Other Communication
Disorders, National Institutes of Health, Bethesda,
MD, USA

Casey E. Reed

National Toxicology Program (NTP) Laboratories
Branch, Division of the NTP, National Institute of
Environmental Health Sciences, National Institutes
of Health, Department of Health and Human Services,
Research Triangle Park, NC, USA

Jenni Repo

School of Pharmacy/Toxicology, Faculty of Health
Sciences, University of Eastern Finland, Kuopio, Finland

Matthew R. Ricci

Research Diets, Inc., New Brunswick, NJ, USA

Anabela P. Rolo

Center for Neurosciences and Cell Biology, University
of Coimbra, Department of Biology, University of
Aveiro, Portugal

Magdalini Sachana

Laboratory of Biochemistry and Toxicology,
Faculty of Veterinary Medicine, Aristotle University
of Thessaloniki, Thessaloniki, Greece

Heidi Sahlman

School of Pharmacy/Toxicology, Faculty of
Health Sciences, University of Eastern Finland,
Kuopio, Finland

Nitin Saini

Janssen Research & Development, Malvern, PA, USA

William F. Salminen

PAREXEL, Sarasota, FL, USA

Kai Savolainen

Nanosafety Research Centre, Finnish Institute of
Occupational Health, Helsinki, Finland

Laura K. Schnackenberg

Division of Systems Biology, National Center
for Toxicological Research, US Food and Drug
Administration, Jefferson, AR, USA

D.T. Selvam

Division of Microbiology, Division of Regulatory
Toxicology, Defence Research and Development,
Gwalior, MP, India

Praveen Sharma

Professor and Head, Department of Biochemistry,
All India Institute of Medical Sciences, Jodhpur, India

Qiang Shi

Division of Systems Biology, National Center for
Toxicological Research, US Food and Drug
Administration, Jefferson, AR, USA

Elina Sieppi

Department of Pharmacology and Toxicology, Institute
of Biomedicine, and Center for Arctic Medicine, Thule
Institute of Oulu, University of Oulu, Oulu, Finland

Jon Sobus

US Environmental Protection Agency, National
Exposure Research Laboratory, Research Triangle Park,
NC, USA

Miguel A. Sogorb

Instituto de Bioingeniería, Universidad Miguel Hernández de Elche, Spain

Melissa Stick

Chief, Scientific Review Branch, Division of Extramural Activities, National Institute on Deafness and Other Communication Disorders, National Institutes of Health, Bethesda, MD, USA

Markus Storvik

School of Pharmacy/Toxicology, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland

David T. Szabo

National Center for Environmental Assessment, Office of Research and Development, US Environmental Protection Agency, Washington, DC, USA

Yu-Mei Tan

US Environmental Protection Agency, National Exposure Research Laboratory, Research Triangle Park, NC, USA

João S. Teodoro

Department of Life Sciences, Center for Neurosciences and Cell Biology, University of Coimbra, Portugal

Rogelio Tornero-Velez

US Environmental Protection Agency, National Exposure Research Laboratory, Research Triangle Park, NC, USA

Beáta Toth

SOLVO Biotechnology, Budaörs, Hungary

Aristides M. Tsatsakis

Department of Forensic Sciences & Toxicology, Faculty of Medicine, University of Crete, Heraklion, Greece

Kirsi Vähäkangas

School of Pharmacy/Toxicology, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland

Deon van der Merwe

Kansas State University, Department of Diagnostic Medicine/Pathobiology, Kansas State Veterinary Diagnostic Laboratory, Manhattan, KS, USA

Ana T. Varela

Department of Life Sciences, Center for Neurosciences and Cell Biology, University of Coimbra, Portugal

Eugenio Vilanova

Instituto de Bioingeniería, Universidad Miguel Hernández de Elche, Spain

Suryanarayana V. Vulimiri

National Center for Environmental Assessment, Environmental Protection Agency (EPA), Washington DC, USA

Kevin D. Welch

United States Department of Agriculture, Agricultural Research Service, Poisonous Plant Research Laboratory, Logan, UT, USA

Xi Yang

Division of Systems Biology, National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR, USA

Snjezana Zaja-Milatovic

School of Medicine, University of Virginia, Charlottesville, VA, USA

Csaba K. Zoltani

Emeritus US Army Research Laboratory, Aberdeen Proving Ground, MD, USA

Introduction

Ramesh C. Gupta

Biomarkers can be broadly defined as indicators or signaling events in biological systems or samples of measurable changes at the molecular, biochemical, cellular, physiological, pathological, or behavioral levels in response to xenobiotics. The Biomarkers Definitions Working Group of the National Institutes of Health (NIH) has defined the biomarker as “a characteristic that is objectively measured and evaluated as an indicator of normal biological processes or pharmacological responses to a therapeutic agent.” In the field of toxicology, biomarkers have been classified as markers of exposure, effect, and susceptibility. Measurement of biomarkers reflects the time-course of an injury and provides information on the molecular mechanisms of toxicity. These biomarkers provide us the confidence of accurate diagnosis, prognosis, and treatment. The biomarkers of early chemical exposure can occur in concert with biomarkers of early disease detection, and that information aids in avoiding further chemical exposure and in strategic development of a novel treatment, including personalized medicine (i.e. treating the patient, and not the disease). In essence, with the utilization of specific biomarkers, an ounce of prevention can be worth a pound of treatment.

Biomarkers are used in drug development, during preclinical and clinical trials, for efficacy and safety assessment. Safety biomarkers can be used to predict, detect, and monitor drug-induced toxicity during both preclinical studies and human clinical trials. Developing highly sensitive methods and their validation for measurement of biomarkers and understanding the resultant data are complex processes that require a great deal of time, effort, and intellectual input. Furthermore, understanding drug metabolism seems essential in

some cases, as the metabolite of a drug can be used as a biomarker, and the drug and/or its metabolite has to be patented by the United States Patent Office and by a similar governmental office/agency in other countries. In the past, many drugs were developed with biomarker assays that guided their use, and this trend is likely to continue in the future for drug discovery and development. With the judicious use of biomarkers, as in evidence-based medicine, patients are most likely to benefit from select treatments and least likely to suffer from their adverse effects. On the contrary, utilization of a bad biomarker can be as harmful to a patient as a bad drug. Therefore, biomarkers need to be validated and evaluated by an accredited laboratory, which participates in a proficiency testing program, to provide a high level of confidence to both clinicians and patients.

In the toxicology field, biomarkers should be specific, accurate, sensitive, valid, biologically or clinically relevant, and easy and fast to perform in order to be useful as predictive tools for toxicity testing and surveillance and for improving quantitative estimates of exposure and dose. Therefore, biomarkers are utilized in biomonitoring data that are useful in a variety of applications, from exposure assessment to risk assessment and management.

In the early 1990s, Dr. Maria Cristina Fossi, from the University of Siena, Italy, emphasized the approach for the development and validation of nondestructive biomarkers over destructive biomarkers in the field of toxicology. She described the ideal biomarker as being: measurable in readily available tissues or biological products and obtainable in a noninvasive way; related to exposure and/or degree of harm to the organism; directly related to the mechanism of action of the contaminants;

highly sensitive with techniques that require minimal quantities of sample and are easy to perform and cost effective; and suitable for different species. The development and validation of new techniques in the laboratory may provide the basis for a valuable field method. But, before a new biomarker's application, some basic information is required, such as dose–response relationships, and biological and environmental factors, which can influence the baseline values of responses. It is important to mention that, when dealing with a biochemical or metabolic biomarker, species differences can be the biggest challenge for any toxicologist.

Biomarkers have applications in all areas of toxicology, especially in the field of pesticides, metals, mycotoxins, and drugs. In the case of veterinary toxicology, biomarkers of plant toxins deserve equal attention. Farmers, pesticide application workers, and greenhouse workers are exposed to pesticides by direct contact and their family members can be exposed via second-hand exposure. Measurement of residues of pesticides and their metabolites, and metals in urine, serves as the most accurate and reliable biomarkers of exposure in agriculture, industrial, and occupational safety and health settings. Recent evidence suggests that *in utero* or early life-exposure to certain pesticides, metals, and other environmental contaminants may cause neurodegenerative (Alzheimer's, Parkinson's, schizophrenia, Huntington's, ALS, and others), metabolic, and cardiovascular diseases, and cancer later in life. In these diseases and many others, specific and sensitive biomarkers play important roles in early diagnosis, and this can serve as the cornerstone for timely therapeutic intervention.

Mycotoxin related toxicity, carcinogenesis, and other health ailments are encountered in man and animals around the world. In developing countries, where regulatory guidelines are not strictly followed, adverse health effects (especially reproductive and developmental effects) are devastating. In these scenarios, early biomarkers of exposure play a pivotal role in avoiding further exposure to the contaminated food/feed and thus safeguard human and animal health.

With the current knowledge of system biology, metabolomics, and various mathematical and computational/chemometric modelings, undetectable biomarkers can be discovered and these biomarkers can predict how tissues respond to toxicants and drugs and/or their metabolites, and how the tissue damage and repair processes compromise the tissue's function. Imaging and chemometric biomarkers are of greater sensitivity and carry more information than conventional biomarkers, as they detect: (1) low levels of chemical exposure (exposure biomarker), and (2) an early tissue response (endogenous response biomarker). The priority will always be for the development of a noninvasive approach over an

invasive approach, and nondestructive biomarkers over destructive biomarkers, but this may not be possible in all cases. Recently, the 2011 Joint SOT/EUROTOX-Debate proposed that “biomarkers from blood and urine will replace traditional histopathological evaluation to determine adverse responses,” identifying and comparing the strengths and limitations of histopathology with serum and urine biomarkers. Unlike histopathological techniques, blood and urine biomarkers are noninvasive, quantifiable, and of translational value. Of course, the complete replacement of histopathological biomarkers with blood and urine may not be possible in the near future, as in some instances histopathological biomarkers will still be used because of recent developments in invaluable molecular pathology techniques.

For the quest of developing the most sensitive and reliable biomarkers, integration of novel and existing biomarkers with a multidisciplinary approach appears fruitful. Furthermore, a multi-biomarkers approach seems more informative and accurate than a single biomarker approach. The latest technologies, such as microRNAs (miRNAs), have been well recognized as reliable and robust biomarkers for early detection of diseases, birth defects, pathological changes, cancer, and toxicities. Because they are stable in biofluids, such as blood, there is rapidly growing interest in using miRNAs as diagnostic, prognostic, and predictive biomarkers, and the outlook for the clinical application of miRNA discoveries is promising, especially in molecular medicine. The application of miRNAs is still very new. Soon incorporating pharmacological and toxicological targeting of miRNAs into the development of innovative therapeutic strategies will be routine. Still, more innovative biomarkers need to be developed that will be highly sensitive (biotechnology-based techniques), require minimum quantities of sample, and will promise high throughput screening.

At the recent annual meetings of the Society of Toxicology in San Francisco, CA, and San Antonio, TX, USA, the EUROTOX in Stockholm, Sweden, and International Congress of Toxicology in Seoul, South Korea, a large number of toxicologists emphasized the importance of biomarkers in health, disease, and toxicity. Accordingly, the *Biomarkers in Toxicology* book is prepared to meet the challenges of today's toxicologists, pharmacologists, and physicians in academia, industry, and government. This reference book may be of particular interest to those in governmental agencies, such as USEPA, USFDA, NIOSH, OSHA, CDC, REACH, EFSA, etc. This is the most comprehensive biomarkers book to date as it covers every possible aspect of exposure, effects, and susceptibility to chemicals. There are many novel topics that are not covered in any previous book. This book identifies and establishes the most sensitive, accurate, unique, and validated biomarkers that can be

used as indicators of exposure and effect(s) of chemicals, and chemical-related long-term diseases, such as cardiovascular, metabolic and neurodegenerative diseases, and cancer. Sixty-four chapters are organized under eight sections with a user-friendly format, and each chapter is enriched with current literature and references for further reading. The book begins with general concepts of toxicity and safety testing and biomarker development using various animal and animal alternative models, pharmacokinetic/PBPK modeling, followed by biomarkers of system/organ toxicity, chemicals and biotoxins. There are several chapters on biomarkers of pharmaceuticals, nutraceuticals, petroleum products, chemical mixtures, radiation, nanoparticles, epigenetics, genotoxicity, and carcinogenesis. Seven chapters are devoted to special topics including biomarkers of neurodegenerative diseases (Alzheimer's and Parkinson's), mitochondrial dysfunction and toxicity, oxidative/nitrosative stress, citrulline as a biomarker in diseases and toxicities, and pathological biomarkers. Lastly, a large number of chapters are

dedicated to the application of biomarkers in toxicology, including the latest strategies and technologies in the development of biomarkers, biomarkers in drug development, safety evaluation, and toxicity testing and integration of biomarkers in biomonitoring of chemical exposure and risk assessment, especially in the context of environmental and occupational medicine.

The editor remains indebted to the contributors of this book for their hard work and dedication. These contributors are highly qualified and considered authorities in the fields of toxicology and pharmacology. He expresses his gratitude to Ms. Denise Gupta for indexing, and to Ms. Robin B. Doss and Ms. Michelle A. Lasher for their continuous support in technical assistance and text and reference checking. Finally, the editor would like to thank Ms. Mary Preap, Ms. Kristine Jones, Ms Marion Stockton, Ms Melissa Read and Mr Roderick Crews (Academic Press/Elsevier) for their immense support at every stage of the production of this book.

Rodents model for toxicity testing and biomarkers

Shayne C. Gad

INTRODUCTION

Three rodent species are widely used in toxicology: the rat, the mouse, and the hamster. Two of these, the rat and mouse, are the most widely used in experimental biology and medicine. These have formed the basis for the identification of toxicities associated with exposure to drugs, industrial and agricultural chemicals, evaluation of their potential hazard, and understanding the mechanisms of their toxicity since toxicology became an identified discipline.

A large set of biomarkers are known for use in identifying and determining the relative (and relevant) risks to humans or other target species. These include:

- Body weights
- Hematology
- Clinical chemistry
- Organ weights
- Gross histologic changes at necropsy
- Microscopic
- Changes in physiologic functions and electrophysiology

Biomarkers are measurements of test model (animal) parameters that can provide important quantitative data about the biological state of the test model, predictive of effects in humans. These biomarkers in toxicology are preferably shared by both test animals and humans,

and in a manner that the relationship of findings in one species as applying to another is known.

Accordingly, we will proceed to understand the current uses of these three rodent species as predictive models for effects in humans, as well as how they are measured and what their normal ranges are.

As these potential pieces of data are overviewed and considered, it is important to remember that each of these biomarkers is a part of the overall picture as to what the model is predicting as per potential adverse effects in humans. Meaningful safety assessment requires that all the data be incorporated in an integrated safety assessment.

The picture becomes both more complex but also clearer as to relevance as new biomarkers are identified and come to be understood. These include proteomics (Amacher, 2010), new clinical chemistry parameters, immune system responses, and real-time physiologic system measurements by telemetrized instrumentation (Gad, 2013).

Table 2.1 (Gad, 2013) presents the classic associations of biomarkers with renal and liver toxicity, while Table 2.2 (adapted from Gad, 2013) presents an overview of the association between classical clinical chemistry parameters and specific target organ toxicities. Table 2.3 summarizes causes associated with hematological findings in the rat.

In the last few years, diligent efforts under the rubric of the Critical Path Initiative have led to the

TABLE 2.1 Classic associations in toxicology

Liver Toxicity	Renal Toxicity
Increased plasma activity of liver marker enzymes: e.g. alanine and aspartate aminotransferases.	Increased water consumption and urine volume. Urine parameters may change, e.g. enzymes and cellular debris.
Decreased plasma total protein concentration.	Increased plasma concentrations of urea and creatinine. Proteinuria.
Increased coagulation times due to decreased synthesis of coagulation factors.	Severe renal toxicity may lead to decreased erythrocyte parameters due to effects on erythrocyte synthesis.
Increased liver weight due to enzyme induction or accumulation of lipid or glycogen.	Increased kidney weight.
Change in color or size at necropsy.	Change in color or size at necropsy.
Histological findings such as necrosis or centrilobular hypertrophy due to enzyme induction.	Histological change, e.g. basophilic tubules or necrosis, papillary necrosis, or glomerular changes.

TABLE 2.2 Association of changes in biochemical parameters with actions at particular target organs

Parameter	Blood	Heart	Lung	Kidney	Liver	Bone	Intestine	Pancreas	Notes
Albumin				↓	↓				Produced by the liver; very significant reductions indicate extensive liver damage
ALP					↑	↑	↑		Elevations usually associated with cholestasis; bone alkaline phosphatase tends to be higher in young animals
ALT (formerly SGPT)					↑				Elevations usually associated with hepatic damage or disease
AST (formerly SGOT)		↑		↑	↑			↑	Present in skeletal muscle and heart and most commonly associated with damage to them
Beta-2-microglobulin				↑					
Bilirubin (total)	↑				↑				Usually elevated due to cholestasis, due to either obstruction or hepatopathy
BUN				↑	↓				Estimates blood filtering capacity of the kidneys; does not become significantly elevated until the kidney function is reduced to 60–75%
Calcium				↑					Can be life threatening and result in acute death
Cholinesterase				↑	↓				Found in plasma, brain, and RBC
CPK		↑							Most often elevated due to skeletal muscle damage but can also be produced by cardiac muscle damage; can be more sensitive than histopathology
Creatinine				↑					Also estimates blood filtering capacity of kidney as BUN does
Glucose								↑	Alterations other than those associated with stress uncommon and reflect an effect on the pancreatic islets or anorexia
GGT					↑				Elevated in cholestasis; this is a microsomal enzyme, and levels often increase in response to microsomal enzyme induction
HBDH		↑			↑				—
KIM-1				↑					
LDH		↑	↑	↑	↑				Increase usually due to skeletal muscle, cardiac muscle, or liver damage; not very specific
Protein (total)				↓	↓				Absolute alterations usually associated with decreased production (liver) or increased loss (kidney); can see increase in case of muscle wasting (catabolism)
SDH					↑↓				Liver enzyme that can be quite sensitive but is fairly unstable; samples should be processed as soon as possible
Troponin		↑							

Note: ↑, increase in chemistry values; ↓, decrease in chemistry values; ALP, alkaline phosphatase; BUN, blood urea nitrogen; CPK, creatinine phosphokinase; GGT, gamma glutamyl transferase; HBDH, hydroxybutyric dehydrogenase; LDH, lactic dehydrogenase; RBC, red blood cells; SDH, sorbitol dehydrogenase; SGOT, serum glutamic oxaloacetic transaminase (also called AST [aspartate amino transferase]); SGPT, serum glutamic pyruvic transaminase (also called ALT [alanine amino transferase]).

TABLE 2.3 Some probable conditions affecting hematological changes

Parameter	Elevation	Depression	Parameter	Elevation	Depression
Red blood cells	<ol style="list-style-type: none"> 1. Vascular shock 2. Excessive diuresis 3. Chronic hypoxia 4. Hyperadrenocorticism 	<ol style="list-style-type: none"> 1. Anemias <ol style="list-style-type: none"> a) Blood Loss b) Hemolysis c) Low RBC production 	Platelets		<ol style="list-style-type: none"> 1. Bone marrow depression 2. Immune disorder
Hematocrit	<ol style="list-style-type: none"> 1. Increased RBC 2. Stress 3. Shock <ol style="list-style-type: none"> a) Trauma b) Surgery 4. Polycythemia 	<ol style="list-style-type: none"> 1. Anemias 2. Pregnancy 3. Excessive hydration 	Neutrophils	<ol style="list-style-type: none"> 1. Acute bacterial infections 2. Tissue necrosis 3. Strenuous exercise 4. Convulsions 5. Tachycardia 6. Acute hemorrhage 	
Hemoglobin	<ol style="list-style-type: none"> 1. Polycythemia (increased in production of RBC) 	<ol style="list-style-type: none"> 1. Anemias 2. Lead Poisonings 	Lymphocytes	<ol style="list-style-type: none"> 1. Leukemia 2. Malnutrition 3. Viral infections 	
Mean cell volume	<ol style="list-style-type: none"> 1. Anemias 2. B-12 deficiency 	<ol style="list-style-type: none"> 1. Iron deficiency 	Monocytes	<ol style="list-style-type: none"> 1. Protozoal infections 	
Mean corpuscular hemoglobin	<ol style="list-style-type: none"> 1. Reticulocytosis 	<ol style="list-style-type: none"> 1. Iron deficiency 	Eosinophils	<ol style="list-style-type: none"> 1. Allergy 2. Irradiation 3. Pernicious anemia 4. Parasitism 	
White blood cells	<ol style="list-style-type: none"> 1. Bacterial infections 2. Bone marrow stimulation 	<ol style="list-style-type: none"> 1. Bone marrow depression 2. Cancer chemotherapy 3. Chemical intoxication 4. Splenic disorders 	Basophils	<ol style="list-style-type: none"> 1. Lead poisoning 	

identification of a more specific set of clinical chemistry biomarkers for key potential target organs, and these can be shown in this grouping:

Heart	Troponins	Zethelius et al. (2008)
Kidney	KIM-1, Albumin, Beta-2-Microglobulin	Hoffmann et al. (2010) ; Ozer et al. (2010) ; Vaidya et al. (2010)
Liver	DILI (Drug Induced Liver Injury), ALT, BUN, coagulation factor	Shi et al. (2010)

Other recently identified biomarkers for specific targets include:

- Mitochondrial Dysfunction.* Increased uptake of calcium (because of ATP depletion) by mitochondria activates phospholipases, resulting in accumulation of free fatty acids. These cause changes in the permeability of mitochondrial membranes, such as the *mitochondrial permeability transition*.
- Progressive Loss of Phospholipids.* Increased degradation by endogenous phospholipases and inability of the cell to keep up with synthesis of new phospholipids (reacylation, an ATP-dependent process).
- Cytoskeletal Abnormalities.* Activated proteases lyse cytoskeletal elements and cell swelling causes detachment

- of cell membrane from cytoskeleton; stretching of the cell membrane results in increased membrane damage.*
- Reactive Oxygen Species.* Produced within the cell and by infiltrating neutrophils and macrophages, especially after restoration of blood flow to an area (reperfusion injury). Cell injury triggers release of a number of inflammatory cytokines and chemokines which amplify the host immune response and attract neutrophils to the site.
- Lipid Breakdown Products.* Unesterified free fatty acids, acyl carnitine, and lysophospholipids. These have a detergent effect on membranes and may exchange with membrane phospholipids, causing permeability changes.

THE RAT

Use in toxicological research

Ideally, safety testing of products intended for use in humans, or to which humans could be exposed, should be done in humans. The data from humans would apply without reservation to complex human physiology and cellular/biochemical mechanisms and human risk assessment. Unfortunately, humans cannot be used

for this purpose. Therefore, the choice of an appropriate species for toxicology studies should be based upon a comparison of the pharmacokinetics and metabolism of the test compound in different laboratory species and man. In the absence of this data this choice is often based upon practicality and economics. The rat has become a species of choice because of metabolic similarities, as well as their small size, relatively docile nature, short life span, and short gestation period. The extensive use of the rat in research has led to the development of a large historical database of their nutrition, diseases, and general biology.

Characteristics

While the rat is a species of choice in toxicology research because of the many physiological similarities and anatomical characteristics, differences exist that must be considered when designing and conducting studies with this animal. Rats are obligate nose breathers, and, as such, an inhaled test material is subject to nasal filtration and absorption. The placenta is considerably more porous in the rat. This difference may increase the chance of fetal exposure to an administered test material or increase the overall level of fetal exposure to an administered test material. The overall distribution of intestinal microflora is different in the rat, which may lead to differences in the metabolism of an orally administered test material. These and other differences in the rat may lead to positive signs of toxicity to a test material that may not be present in a different species.

Strain differences

Breeding rats for specific characteristics has produced some physiological differences between strains of rats. Some of these differences are known to affect how the various strains react to toxicants. Among others, strain-specific differences have been found in sensitivity to thiourea (Dieke and Richter, 1945), sensitivity to acetaminophen nephrotoxicity (Newton *et al.*, 1985a,b), the

incidence of spontaneous glomerular sclerosis (Bolton *et al.*, 1976), sensitivity to the carcinogenic actions of 7,12-dimethylbenz(a)anthracene (Boyland and Sydnor, 1962), the effects of trimethyltin on operant behavior and hippocampal GFAP (MacPhail *et al.*, 2003), differences in renal carcinogenesis (Hino *et al.*, 2003), differences in cytochrome P4501A1 gene expression caused by 2,3,7,8-tetrachlorodibenzo-p-dioxin in the liver (Jana *et al.*, 1998), susceptibility to 4-nitroquinoline 1-oxide inducing carcinoma (Kitano *et al.*, 1992), and differences in the levels of drug-metabolizing enzymes (Page and Vesell, 1969). In recent years, research and breeding programs have been focused on producing inbred and outbred strains focused on specific disease models and susceptibility to the development of certain carcinoma. When choosing a strain for use, it is important to consider these differences.

Of importance for carcinogenicity studies, strain differences have been found in the incidence of spontaneous tumors. Table 2.4 gives the incidence of spontaneous tumors found in commonly used strains in carcinogenicity studies. The historical incidence is important to the analysis of a study in that a high spontaneous rate may mask a small test material related increase in tumor incidence.

Due to lower spontaneous tumor rates, the Wistar has become the most popular strain in toxicological research.

Normal physiological values

General values for selected physiological parameters are given in Tables 2.5 and 2.6. Normal values will vary based upon the strain of animal, supplier, feed, and housing conditions. These tables should be used as a point of reference only.

Study designs

The length and design of toxicology studies used to predict human risk are governed by guidelines issued by

TABLE 2.4 Incidence of common spontaneous tumors in Fischer 344 and CD (SD)IGS rats

		% Tumors in Untreated Rats							
Organ	Tumor Type	CD(SD)IGS		CD (SD)		Fisher		Wistar	
		Male	Female	Male	Female	Male	Female	Male	Female
Adrenal gland	Pheochromocytoma	10.0	2.3	11.3	2.3	11.9	3.2	3.2	1.3
Mammary gland	Fibroadenoma	1.4	44.5	1.3	16.7	0.8	7.1	1.2	30.2
Pancreas	Islet cell adenoma	3.6	1.4	4.0	0.3	1.5	0.2	5.3	1.9
Pituitary Gland	Adenoma pars distalis	33.6	56.8	35.7	50.3	12.4	28.2	41.1	65.8
Testis	Interstitial cell tumor	1.8		7.0		74.6		4.3	
Thyroid gland	C-Cell Adenoma	10.5	5.0	5.0	5.7	12.5	8.2	10.1	10.7

Source: Adapted from Charles River (2000), Mitsumori *et al.* (2001)

TABLE 2.5 Selected normative data

Husbandry	
Room temperature (°C)	18–26
Relative humidity (%)	30–70
Ventilation (air change/h)	10
Light/dark cycle (hours)	12–14/12–10
Minimum cage floor size	
Housed individually (cm ²)	350
Breeding with pup (cm ²)	800
Group housed (cm ² adult)	250
General	
Lifespan (years)	2.5–3.0
Surface area (cm ²)	0.03–0.06
Chromosome number (diploid)	42
Water consumption (mL/100g/day)	10–12
Food consumption (g/day)	20–40
Average body temperature (°C)	37.5
Reproduction	
Puberty (males and females)	50 ± 10 days
Breeding season	All year
Type of estrous cycle	Polyestrous
Length of estrous cycle	4–5 days
Duration of estrous	10–20 hr
Mechanism of ovulation	Spontaneous
Time of ovulation	7–10 hr after onset of estrus
Time of implantation	Late day 4 or 5 ^a
Length of gestation	21–23 days
Litter size	8–16 pups
Birth weight	5–6 g
Eyes open	10–12 days
Weaning age/weight	21 days/40–50 g
Cardiovascular	
Arterial blood pressure	
Systolic (mmHg)	116–145
Diastolic (mmHg)	76–97
Heart rate (beats/min)	296–388
Cardiac output (mL/min)	10–80
Blood volume (mL/kg)	64
Pulmonary	
Respiration (Breaths/min)	100–140
Tidal Volume (mL)	1.1–2.5
Compliance (mL/cm H ₂ O)	0.3–0.9
Resistance (cm H ₂ O/mL*s)	0.1–0.55
Pattern	Obligate nasal
Renal	
Urine volume	15–30 mL/24 hour
Na ⁺ excretion	200 mmol/L/24 hour
K ⁺ excretion	150 mmol/L/24 hour
Urine osmolality	2000 mOsm/kg H ₂ O
Urine pH	7.3–8.5
Urine Specific Gravity	1.01–1.07
Urine Creatinine	6 μmol/L/24 hour
Glomerular Filtration Rate	1.0 mL/min/100 g body weight

Source: Data from Peplow *et al.* (1974), Baker *et al.* (1979), Bivin *et al.* (1979), Waynforth and Flecknell (1980), Van Zutphen *et al.* (1993), and Sharp and La Regina (1998).

^aThe estrous cycle length may vary from 4–5 days between strains. Time of implantation may vary based upon the length of the estrous cycle and is dependent upon Day 0 or the first day sperm is found in the vagina.

regulatory bodies such as the United States Food and Drug Administration (FDA), the International Council on Harmonization (ICH), the Environmental Protection Agency (EPA), and their counterparts worldwide. Toxicology studies are divided into a series of three sets of studies that are required for each phase of clinical trials. For initial approval to begin clinical trials, the following studies are required. The length of dosing in the toxicology studies varies depending upon the intended length in clinical trials. A test compound intended to be a repeat dose study for up to 28 days in duration initially requires a two-phase study in which a maximum tolerated dose (MTD) following a single administration is determined followed by a second phase during which the test compound is administered daily at dose levels based upon the MTD for 5 to 7 days (Table 2.7). Following the completion of the MTD study a 14 or 28 Day Repeat Dose study (Table 2.8) or a 28 Day Repeat Dose study with immunophenotyping (Table 2.9) should be conducted. These studies assess the effects of a test compound at dosages that do not cause immediate toxic effects.

In support of Phase II clinical trials, longer-term subchronic and chronic toxicity studies (Table 2.10) should be conducted. Subchronic and chronic toxicity studies are designed to assess the test compound effects following prolonged periods of exposure. The highest dosage level in each of these studies should produce a toxic effect such that target organ may be identified. The lowest dosage level should provide a margin of safety that exceeds the human clinical dose and ideally allows for the definition of a no observable effect level (NOEL). Alternatively, when effects related to the pharmacological mechanism of the test compound, or when observed effects may be related to treatment with the test compound but may not be of toxicologic significance, a no observable adverse effect level (NOAEL) may be determined.

In addition to the subchronic and chronic toxicity studies in support of Phase II clinical trials, reproductive safety studies may also be required. Reproductive toxicity studies are typically required for test compounds intended to be administered to women of child-bearing age or which may affect male reproduction. These studies include an assessment of the potential effects of the test compound on general fertility and reproductive performance (Segment I), developmental toxicity (Segment II), or perinatal and postnatal development (Segment III). The highest dose in reproductive studies should be chosen so that administration causes some minimal toxicity. Typically, a dose range finding pilot study in a small number of animals should be conducted prior to initiating the definitive reproductive toxicology studies. Examples of protocols designed to meet the ICH guidelines are presented in Tables 2.11, 2.12, and 2.13.

TABLE 2.6 Growth rates in selected rat strains

Weight (g)	Age (days)							
	CrI:CD (SD)IGSBR		CrI:(WI)BR		CrI:(LE)BR		CDF(F-344)/CrIBR	
	Males	Females	Males	Females	Males	Females	Males	Females
Up to 50	Up to 23	Up to 23	Up to 23	Up to 25	Up to 21	Up to 21	Up to 23	Up to 23
51–75	24–28	24–29	24–28	26–30	22–25	22–26	24–29	24–29
76–100	29–34	30–35	29–32	31–34	26–29	27–31	30–34	30–35
101–125	35–37	36–39	33–35	35–40	30–34	32–36	35–39	36–42
126–150	38–42	40–44	36–40	41–47	35–37	37–43	40–45	43–55
151–175	43–45	45–50	41–44	48–56	38–42	44–50	46–50	56–72
176–200	46–49	51–56	45–48	57–64	43–46	51–55	51–57	73–105
201–225	50–52	57–70	49–52	65–81	47–49	56–69	58–63	105 +
226–250	53–56	71–84	53–56	82–105	50–55	70–86	64 +	
251–275	57–59	84–105	57–61	106 +	56–58	87–102		
276–300	60–65	106 +	62–67		59–64	103 +		
301–325	66–71		68–73		65–70			
326–350	72–77		74–79		71–80			
351–375	78–87		80–87		81–90			
376 +	88 +		88 +		91 +			

Source: Adapted from Charles River (2000)

TABLE 2.7 Maximum tolerated dose study in rats

Phase A	Oral MTD Study	
	Males	Females
Dose Level 1	3	3
Dose Level 2	3	3
Dose Level 3	3	3
Dose Level 4	3	3

Phase B	7-Day Oral Range Finding Study			
	Main Study		Toxicokinetics	
	Males	Females	Males	Females
Control	5	5	–	–
Low Dose	5	5	9	9
Mid Dose	5	5	9	9
High Dose	5	5	9	9

EXPERIMENTAL DESIGN:

In Phase A, the dose level will be increased until the maximum tolerated dose (MTD) is determined. The MTD is a dose that produces neither mortality nor more than a 10% decrement in body weight nor clinical signs of toxicity. In Phase B, animals will be dosed daily for 7 days at fractions of the single dose MTD to estimate a repeat dose MTD.

Dose Route/Frequency:

As requested

Phase A: Once

Phase B: Once per day for 7 consecutive days

OBSERVATIONS: Twice daily in both Phases (mortality/morbidity)

DETAILED CLINICAL OBSERVATIONS: Daily in both phases

BODY WEIGHTS: Daily in both phases

FOOD CONSUMPTION: Daily

CLINICAL PATHOLOGY (Phase B only): Hematology, clinical chemistry, and urinalysis evaluations on all surviving main study animals at termination.

NECROPSY (Phase B only): Tissues saved for possible future histopathological evaluation.

In support of Phase III clinical trials, a carcinogenicity study may be required (Table 2.14). Typically 18 months to 2 years in duration, this type of study is designed to assess the potential of the test compound to induce neoplastic lesions. The highest dosage in a carcinogenicity study should cause minimal toxicity when administered via the intended route for clinical use. The preclinical studies required in support of the clinical trials are dependent upon the intended route and frequency of administration of the test compound and the intended age group to be treated.

Routes of test article administration

Oral routes

Rodents have several unique characteristics to be considered regarding the oral administration of test compounds. One of the most important characteristics is the lack of an emetic response. The lack of this response allows for a higher dose of a potential emetic compound to be administered and evaluated. Many compounds and excipients may cause emesis in dogs or other large animal species and may lead to a low level of exposure and erratic blood levels. A second factor to consider is

ORGAN WEIGHTS (Phase B only): Adrenals, brain, heart, kidneys, liver, lungs, ovaries with oviducts, pituitary, prostate, salivary glands, seminal vesicles, spleen, thyroid with parathyroid, thymus, testes, uterus.

TOXICOKINETICS: Blood collected on Days 1 and 7 (3 cohorts consisting of 3 animals/sex/treatment group bled twice to equal six time points), calculation of C_{max} , T_{max} , AUC_{0-24} , and $T_{1/2}$.

TABLE 2.8 14 or 28 day repeat dose toxicity study in rats

	Main Study		Toxicokinetics	
	Males	Females	Males	Females
Vehicle Control	10	10	—	—
Low Dose	10	10	9 + 3**	9 + 3**
Mid Dose	10	10	9 + 3**	9 + 3**
High Dose	10	10	9 + 3**	9 + 3**

**Three additional animals/sex/treatment group included as replacement animals.

OBSERVATIONS: Twice daily (mortality/morbidity)

DETAILED CLINICAL OBSERVATION: Weekly

FUNCTIONAL OBSERVATIONAL BATTERY: Pretest and Day 14 or 25

BODY WEIGHTS: Weekly

FOOD CONSUMPTION: Weekly

OPHTHALMOLOGY: All animals prior to test article administration; all surviving main study animals at study termination.

CLINICAL PATHOLOGY: Hematology, clinical chemistry, and urinalysis evaluations on all surviving main study animals at termination.

TOXICOKINETICS: Blood collected on Days 1 and 14 or 27 (3 cohorts consisting of 3 animals/sex/treatment group bled twice to equal six time points); TK modeling.

NECROPSY: All main study animals; toxicokinetic animals euthanized and discarded.

ORGAN WEIGHTS: Adrenals, brain, heart, kidneys, liver, lungs, ovaries with oviducts, pituitary, prostate, salivary glands, seminal vesicles, spleen, thyroid with parathyroid, thymus, testes, uterus.

SLIDE PREPARATION/MICROSCOPIC PATHOLOGY: All animals in the vehicle control and high dose groups and all found dead animals: full set of standard tissues; low and mid dose group target organs (to be determined); gross lesions from all animals.

that rodents are nocturnal and eat most of their food at night. When maintained on a 12-hour light-dark cycle, rats have been found to consume 75% of their daily food intake during the dark cycle. This should be taken into consideration when designing an oral gavage study and determining when the animal may be dosed. Early in the light cycle, animals are more likely to have a full stomach and complications associated with dosing may occur if large volumes of test article are administered. In addition, a full stomach may affect gastric emptying and the rate of absorption of an orally administered test compound.

Techniques for oral administration of test compounds include mixing in the diet, via gavage or stomach tube, via capsule, or in drinking water. The most widely used methods of oral administration are the dietary and gavage techniques.

Dietary vs. gavage methods

The choice between dietary and gavage dosing techniques is typically based upon several factors. A scientific decision can only be made with a knowledge of the pharmacokinetics of the test compound administered

TABLE 2.9 28 day repeat dose toxicity study with immunophenotyping in rats

	Main Study		Toxicokinetics	
	Males	Females	Males	Females
Vehicle Control	10	10	—	—
Low Dose	10	10	9 + 3*	9 + 3*
Mid Dose	10	10	9 + 3*	9 + 3*
High Dose	10	10	9 + 3*	9 + 3*

*Three additional animals/sex/treatment groups included as replacement animals; the control animals will not be evaluated for toxicokinetics.

OBSERVATIONS: Twice daily (mortality/morbidity)

DETAILED CLINICAL OBSERVATION: Weekly

FUNCTIONAL OBSERVATIONAL BATTERY: Pretest and Day 25

BODY WEIGHTS: Weekly

FOOD CONSUMPTION: Weekly

OPHTHALMOLOGY: All animals prior to test article administration; all surviving main study animals at study termination.

CLINICAL PATHOLOGY: Hematology, clinical chemistry, and urinalysis evaluations on all surviving main study animals at termination.

IMMUNOTOXICOLOGY: Immunophenotyping of blood leukocytes by flow cytometry on all surviving main study animals at termination. NK cell assay on blood leukocytes of all surviving main study animals at termination.

TOXICOKINETICS: Blood collected on Days 1 and 27 (three cohorts consisting of three animals/sex/treatment group bled twice to equal six time points).

NECROPSY: All main study animals; toxicokinetic animals euthanized and discarded.

ORGAN WEIGHTS: Adrenals, brain, heart, kidneys, liver, lungs, ovaries with oviducts, pituitary, prostate, salivary glands, seminal vesicles, spleen, thyroid with parathyroid, thymus, testes, uterus, two lymph nodes (e.g. mesenteric, axillary, popliteal, etc.) including the lymph node draining the route of administration.

SLIDE PREPARATION/MICROSCOPIC PATHOLOGY: All animals in the vehicle control and high dose groups and all found dead animals: full set of standard tissues (add Peyer's patch, extra lymph node); low and mid dose group target organs; gross lesions from all animals.

by both methods. Other considerations which may be used in making this decision are as follows.

The dietary method can be used with a compound that can be mixed with the diet, is stable under storage conditions in the diet, and is palatable to the animal. A major advantage of the dietary method is that it requires less manpower to perform the study. The diet mixing process can be performed weekly or, if stability allows, less often. The mixing and feeding process is less labor intensive than gavaging rats on a daily basis. Several disadvantages also exist in using the dietary method. Methods must be developed and validated to prove homogeneity and stability. This is not as easy a process as with a suspension or solution. The dietary method is also less exact than the gavage method, in that the concentration of compound mixed in the feed is based upon predicted feed consumption and body weights. In addition, if the feed is not palatable to the animal, or the test compound makes the animal ill, feed

TABLE 2.10 Subchronic and chronic toxicity study in rats

	Main Study		Toxicokinetics	
	Males	Females	Males	Females
Vehicle Control	15	15	—	—
Low Dose	15	15	9 + 3*	9 + 3*
Mid Dose	15	15	9 + 3*	9 + 3*
High Dose	15	15	9 + 3*	9 + 3*

*Three additional animals/sex/treatment group included as replacement animals.

OBSERVATIONS: Twice daily (mortality/morbidity)

DETAILED CLINICAL OBSERVATION: Weekly

BODY WEIGHTS: Weekly

FOOD CONSUMPTION: Weekly

OPHTHALMOLOGY: All animals prior to test article administration; all surviving main study animals at study termination.

CLINICAL PATHOLOGY: Hematology, clinical chemistry, and urinalysis evaluations on all surviving main study animals at termination.

TOXICOKINETICS: Blood collected on Days 1 and 90 (3 cohorts consisting of 3 animals/sex/treatment group bled twice to equal six time points); TK modeling.

NECROPSY: All main study animals; toxicokinetic animals euthanized and discarded.

ORGAN WEIGHTS: Adrenals, brain, heart, kidneys, liver, lungs, ovaries with oviducts, pituitary, prostate, salivary glands, seminal vesicles, spleen, thyroid with parathyroid, thymus, testes, uterus.

SLIDE PREPARATION/MICROSCOPIC PATHOLOGY: All animals in the vehicle control and high dose groups and all found dead animals: full set of standard tissues; low and mid dose group target organs; gross lesions from all animals.

TABLE 2.11 Study of fertility and early embryonic development to implantation in rats

	Males	Females
Vehicle Control	25	25
Low Dose	25	25
Mid Dose	25	25
High Dose	25	25

DOSE ROUTE/FREQUENCY: Males dosed beginning 28 days before mating and continuing until euthanasia. Females dosed beginning 14 days before mating and continuing through Day 7 of gestation (implantation).

OBSERVATIONS: Twice daily (mortality/morbidity)

CLINICAL EXAMINATIONS: Observations for clinical signs, body weights, and food consumption measurements recorded during the study period. Beginning at initiation of test article administration, females examined daily to establish estrous cycle.

UTERINE EXAMINATIONS: Performed on dams on Day 13 of gestation. Gravid uterine weight and the weight of the ovaries recorded. Total number of corpora lutea and implantations, location of resorptions, and embryos recorded. Females subjected to necropsy, and reproductive organs and gross lesions fixed for possible microscopic evaluation.

EVALUATION OF MALES: Following disposition of females, the males euthanized and subjected to a necropsy. The testes and epididymides weighed, and analysis of sperm parameters (concentration, motility, and morphology) performed. Reproductive organs and gross lesions fixed for possible microscopic evaluation.

STATISTICAL ANALYSIS: Standard

TABLE 2.12 Embryo-fetal development in rats

	Time Mated Females
Vehicle Control	25
Low Dose	25
Mid Dose	25
High Dose	25

DOSE ROUTE/FREQUENCY: Dosing initiated on Day 6 of gestation and continuing to and including Day 17 of gestation. **OBSERVATIONS:** Twice daily (mortality/morbidity); **CLINICAL EXAMINATIONS:** Daily, Gestation Days 6 through 20.

BODYWEIGHTS/FOOD CONSUMPTION: Gestation Days 0, 6, 9, 12, 15, 18, and 20. **CESAREAN SECTION/NECROPSY:** Litters will be delivered by cesarean section on Day 20 of gestation. Gravid uterine weight will be recorded. Total number of corpora lutea, implantations, early and late resorptions, live and dead fetuses, and sex and individual body weights of fetuses will be recorded. External abnormalities of fetuses will be recorded. Approximately one-half of the fetuses will be processed for visceral abnormalities, and the remaining fetuses will be processed for skeletal abnormalities. All fetuses will be examined for visceral and skeletal abnormalities. Dams will be subjected to a necropsy and gross lesions and target organs (if known) will be saved.

TABLE 2.13 Pre-and postnatal development, including maternal function in rats

	P Generation		F1 Generation	
	Males	Females	Males	Females
Vehicle Control	NA	25	25	25
Low Dose	NA	25	25	25
Mid Dose	NA	25	25	25
High Dose	NA	25	25	25

NUMBER ON STUDY: P Generation: 100 females, F1 Generation: 100 males, 100 females.

DOSE ROUTE/FREQUENCY: Once daily to P animals from Gestation Day (GD) 6 to Postnatal Day (PND) 21. F₁ animals not dosed.

OBSERVATIONS: Twice daily (mortality/morbidity)

CLINICAL OBSERVATIONS: P females – daily during treatment / F1 adults – weekly

BODY WEIGHTS:

P females – GD 0, 6, 10, 14, 17, and 20, PND 0, 7, 10, 14, and 21

F1 males – weekly through termination

F1 females – weekly until evidence of copulation detected, then GD 0, 7, 10, and 13

FOOD CONSUMPTION: P females – On corresponding body weight days during gestation/lactation

VAGINAL SMEARS: All F1 females during a 21-day cohabitation period until evidence of copulation is detected.

LITTER EVALUATIONS: All F1 offspring, count, body weight, sex, clinical observations on PND 0, 4, 7, 14, 21; behavioral and developmental evaluation of four males and four females from each litter for static righting, pinna detachment, cliff aversion, eye opening, air drop righting reflex, neuropharmacological evaluation, auditory response. One male and one female (selected for the next generation) tested for sexual maturation (vaginal opening, preputial separation), motor activity/emotionality, and passive avoidance.

CESAREAN SECTION: On GD 13, F1 females for location of viable and nonviable embryos, early and late resorptions, number of total implantations, and corpora lutea.

SPERM EVALUATION: May be conducted on F1 males if evidence of reduced fertility is noted (additional cost).

NECROPSY: Gross lesions/target organs fixed for possible microscopic evaluation (additional cost).

All P females at PND 22 as well as all F1 weanlings not selected for F1 generation

All F1 females at GD 13

All F1 males after termination of F1 cesarean sections.

TABLE 2.14 Carcinogenicity study in rats

	Main Study		6-Month Satellite	
	Males	Females	Males	Females
Vehicle Control	60	60	20	20
Low Dose	60	60	20	20
Mid Dose	60	60	20	20
High Dose	60	60	20	20

OBSERVATIONS: Twice daily (mortality/morbidity).

DETAILED CLINICAL OBSERVATIONS: Once weekly.

BODY WEIGHTS: Weekly for first 13 weeks, monthly thereafter.

FOOD CONSUMPTION: Weekly for first 13 weeks, monthly thereafter.

OPHTHALMOLOGY: All animals pretest and all survivors prior to terminal sacrifice.

CLINICAL PATHOLOGY: Main Study: Hematology at termination 6-Month Satellite: Hematology, clinical chemistry, and urinalysis evaluations on all surviving satellite animals at termination.

NECROPSY: All animals.

SLIDE PREPARATION/MICROSCOPIC PATHOLOGY: All animals, full set of standard tissues, all masses and all lesions.

STATISTICAL ANALYSIS: Standard.

consumption may be reduced, thereby reducing exposure to the test compound. In addition, the facility and control animals may be exposed to the test compound through dust or vapors.

The gavage method can be used when the test compound is not stable in the diet or may not be palatable to the animals. In addition, the gavage method is preferable when evaluating toxicokinetics or pharmacokinetics. As with dietary mixtures, test compound administered via gavage as a solution or suspension should be analyzed for homogeneity, stability, and concentration. Methods for solution or suspension may be easier to develop than those required for dietary mixtures. For GLP studies, evaluation of homogeneity, stability, and concentration should be conducted for every study. If the same methodology and batch size are used for multiple studies, homogeneity may be established one time. Stability of the test compound in solution or suspension should be determined under the testing conditions in the proposed vehicle. Typically, stability for toxicology studies is established for between 7 and 14 days. If the test compound is not found to be stable, stability of shorter duration may be established. Lastly, concentration analysis should be established for each dose level and should be periodically evaluated during longer term studies.

With the gavage method of dosing, a more precise amount of the test compound can be delivered and may reduce the amount of test compound required to complete the study. This becomes important when evaluating the effects of a pharmaceutical, as the required dose levels and exposure levels to show safety may be lower than required for a pesticide or chemical. A disadvantage of the gavage method is that it involves handling

of the rat for each dosing. Handling of the rat has been shown to increase corticosterone levels (Barrett and Stockham, 1963) and may affect study results. Additionally, daily intubation may lead to death due to esophageal puncture or inhalation pneumonia.

Dietary method

When utilizing the dietary method, the test compound is mixed with the diet and administered to the animals either *ad libitum* or the diet is presented to the animals for a fixed amount of time each day. The dosage received by an animal is regulated by varying the concentration of test compound in the diet based on the predicted food consumption and body weight. Food consumption and body weight predictions are based upon historical laboratory data for early time points in a study. As the study progresses, growth and food consumption curves can be established for each group and group mean data can be used to predict future food consumption. Different concentrations of the test compound and diet should be made for each sex.

Test compounds and diets are mixed in two steps: (1) the compound and about 10% of the total amount of diet are blended in a premix; then (2) the premix and the remainders of the diet are mixed. The total amount of diet to be mixed is first weighed out, and the 10% is separated into the premix. To make the premix, the entire test compound and an aliquot of the diet (from the 10%) are put into a mortar. These ingredients are ground with a pestle until the mixture appears homogeneous. The mixture and the remainder of the premix are then layered in a small capacity mixer and mixed for 5–10 minutes. The time for this mixing process can be varied if analysis shows the total mixture is not homogeneous. For the final mix, the premix and the remainder of the diet are layered in a large capacity mixer. The mixing time will vary with the type of blender and can be varied if the analysis shows the total mixture is not homogeneous.

Several types of blenders are available for the mixing process; these include open-bowl "kitchen" mixers, V or PK blenders, and Turbula mixers. Metal parts should be ground to eliminate electrostatic forces. In addition, alternative methods of dietary administration such as microencapsulation may be used for volatile, reactive, or unpalatable chemicals.

Gavage method

In the gavage procedure, the test compound is administered by passing a feeding tube or gavage needle attached to a syringe down the esophagus into the stomach.

Test article preparation

If not already a liquid, the test compound is prepared for administration by adding it to the appropriate vehicle. The choice of vehicle will depend on the characteristics of the compound and whether it is to be administered as a suspension or a solution. In addition, consideration must be given to the effects of the vehicle on the rat (Gad and Chengelis, 1998). Common vehicles used include water and food grade oils such as corn oil. Suspensions are made when aqueous vehicles are desired and the test compound is not soluble. Suspending agents such as methylcellulose are added to increase the viscosity and hold the compound in suspension. Other agents such as Tween 80, ethanol, polyethylene glycol 400 (PEG 400), and others may be used as wetting or stabilizing agents.

Equipment

Soft catheters made of silastic or polyethylene (e.g. infant feeding tubes), stainless steel gavage needles with smooth ball-shaped tips, or polyethylene gavage needles with ball-shaped tips are commonly used. All are commercially available and are relatively inexpensive. While the soft catheter minimizes the chance of esophageal trauma, liquid can leak past the catheter and back up the esophagus and be aspirated. The ball-shaped tips of the gavage stainless steel gavage needles reduce the chances of tracheal injections; however, if an animal struggles while the needle is in the esophagus, the rigid needle increases the chances of perforating the esophagus. The polyethylene gavage needle incorporates the best of both the soft catheter and the stainless steel needle, but due to the flexible nature of the needle, the risk for tracheal injection is increased.

Conybeare and Leslie (1988) found deaths in gavage studies were a result of aspiration of small amounts of irritant solutions or acidic, hypertonic solutions. They also found that the use of a ball-tip 4 mm in diameter helped to eliminate deaths related to dosing. With gentle handling, the animals will be acclimated to the techniques used and dosing will become easier.

Aspiration and tracheal administration of test compound as well as esophageal trauma have been associated with gavage dosing and may lead to difficulty in interpretation of the study. The catheter and the needles all have risks inherent in their use, and therefore care should be used when using these tools and animal technicians should be properly trained. The choice of the appropriate catheter or needle should be left up to the technician dosing and should be whatever the technician has been trained in and is most comfortable with.

Technique

The description following is appropriate for either a gavage needle or catheter; for simplicity, only the

needle will be mentioned in the description. Prior to picking up the animal, the syringe should be attached to the needle and filled with the appropriate amount of test compound to be delivered. Any air bubbles should be eliminated and the needle wiped clean of residual test compound. This is done so that the animal does not taste the test compound and residual test compound is not aspirated as the needle is passed down the esophagus. If the dosing liquid is distasteful, the animal may struggle after repeated dosing and increase the chances of being injured.

To position the animal for gavage, it should be grasped by the skin of the back and neck ensuring that the head, neck, and back are in a straight line. Alternatively, the animals can be grasped about the shoulders, with the index finger and thumb on either side of the head. The objective is to firmly hold the animals to be able to control any struggling if it occurs and to also prevent the animal from being able to bite the technician. For even more control, the animal may be placed on a table or brought up against the operator's chest.

Once the animal is in position, the needle can be inserted into the mouth of the animal, and moved over the tongue and down into the esophagus. The length of the needle should be inserted into the animal. A slight rotation of the needle may help with insertion into the esophagus. If the needle is inserted into the trachea, the animal may struggle. The syringe should be grasped lightly such that, if the animal does struggle, the chances of an esophageal tear are minimized. If the animal continues to struggle, the needle should be withdrawn to allow the animal to calm down, and then dosing should be attempted again. Alternatively, if a catheter is used, as the tube is placed into the mouth, it should be placed to the side between the molars. This is done because the tube may be bitten or transected if passed too close to the front teeth.

With the needle in place, the test compound should be slowly expelled into the animal. If administered too rapidly, reflux may occur and the test compound may back up into the esophagus, resulting in an inaccurate dose being given and possible aspiration of the test compound. Once the dose has been delivered, the needle should be withdrawn and the animal observed for any signs of distress or respiratory difficulty. An experienced technician should be able to dose between 5 and 7 animals per minute without causing discomfort to the animals and with minimal dosing-related deaths.

Gavage liquids are commonly administered at a volume of 5–10 mL/kg body weight. The volume should be enough to be delivered accurately, but not so much that it will adversely affect the animal. The maximum volume should be no more than 20 mL/kg. If using volumes greater than 10 mL/kg, it may be advisable to

fast the animals for several hours prior to dosing. This will ensure that the stomach is empty prior to dosing and able to handle the larger volume. This option should be considered carefully, as fasting can affect the rate of absorption and clearance from the stomach. In addition, the choice of housing and bedding should be considered when dosing with large volume, as rats have the tendency to eat the bedding, which may hinder gavage dosing. In addition, the volume chosen can have an effect upon the results of the study and volumes greater than 10 mL/kg should only be used when issues of solubility and exposure exist. Ferguson (1962) found that a change in dose volume of from 5 to 1% of body weight could reduce mortality rate from approximately 95% to 5%, respectively, at equivalent doses.

Neonatal administration

Neonatal intragastric injections can be made orally with thin silicone tubing (Gibson and Becker, 1967) or by intragastric injection with a 27-gauge needle through the abdominal wall (Worth *et al.*, 1963; Bader and Klinger, 1974). The oral method using silicone tubing is performed in a similar manner to the previously described method in adult rats. The intragastric injection through the abdominal wall is performed by first locating the stomach in the upper left quadrant of the abdomen and then carefully inserting the needle through the abdominal wall into the stomach, taking care that the animal does not move. The syringe should be gently aspirated to ensure proper placement and then the injection completed and the needle withdrawn.

Capsule

To eliminate the possibility of dosing errors and to deal with compounds that cannot be delivered through conventional means, methods have been developed for the administration of capsules into the esophagus of the rat. The test compound may be prefabricated into a small capsule or the test article may be weighed and placed into commercially available capsules. An individual capsule is then placed into a specially designed cup in the end of a gavage needle and the needle is then inserted into the esophagus of the rat. The capsule is then pushed out of the cup into the esophagus using either air or a rod inside the needle. The needle is then withdrawn and the capsule moves down into the stomach by peristaltic action. Only a small amount of test compound can be administered as a single dose using this method, but multiple capsules can be administered sequentially in the same dosing session.

Water

As an alternative to dietary administration, compounds that are water soluble, palatable to the rat, and stable in

water may be administered via the drinking water. This method offers similar advantages as adding a test compound to the diet. Additionally, compounds will be more easily mixed and analyses will be more easily developed than when a compound is in the diet. However, spillage of water makes measurement of the actual dose received difficult.

Intravenous route

One of the most common methods of administration of test compound is via intravenous injection or infusion. The intravenous route is often the route of choice for compounds that have poor bioavailability via the oral route or have a short half-life. Several issues must be considered when administering a test compound intravenously. The compound must be soluble in an acceptable intravenous vehicle or excipient, must be able to be administered as a solution, and should be sterile or sterile filtered prior to administration. In addition, when designing a study, the pharmacokinetic profile of the test compound administered intravenously should be considered. Study activities such as clinical observations and functional observational battery should be planned around the expected time of greatest plasma concentration.

A variety of veins may be used for intravenous injections. These include the lateral tail (caudal), jugular, femoral, saphenous, lateral marginal, dorsal metatarsal, sublingual, and dorsal penile vein. While most of these are superficial, and easily available for injection, several require the use of anesthesia or more than one technician and may be of limited use in repeat dose studies. While anesthesia may be acceptable for acute studies or surgical model, its repeated use may have an effect on the toxicity of a test compound.

Lateral tail vein

The lateral tail veins are currently the most widely used for intravenous injections in the rat. The veins are easily visible, especially in young animals, and injections can be performed by one person without the use of anesthesia. The technician performing the function should be well trained and care should be taken to insure that the lateral veins are being accessed and not the dorsal or ventral artery of the vein.

Bolus injection

The animal should be placed in an appropriate restrainer. This typically consists of a solid tube that the animal is placed into head first that has a stop that is placed behind the animal with a hole that allows the tail to hang out the back. The restraint tube is designed to be secure enough that the animal cannot move, back out, or turn, but can still breath comfortably. Once

secure, the tail should be cleaned and the vein may be dilated with heat. This can be accomplished by placing the tail in warm water (40–45°C), placed under a heat lamp, or wrapped with warm gauze. Care must be taken to avoid using excessive heat as tissue damage may result. [Minasian \(1980\)](#) describes a tourniquet made from a plastic syringe and thread. If used, this should not be left on for an extended period of time.

When performing an injection, the end of the tail should be held firmly and taut with the thumb and index finger of one hand. A 23-gauge needle attached to an appropriately sized syringe should be held with the bevel up at a shallow angle parallel to the vein. The skin of the tail is then pierced and the needle advanced until resistance is no longer felt. The plunger of the syringe should then be aspirated to ensure proper placement of the needle. The use of a needle with a clear or transparent hub will facilitate confirmation of correct placement. Blood backflow into the needle confirms entry into the vein. Alternatively, a butterfly needle with an extension line may be used. The butterfly needle with an extension set precludes the need to hold the tail, needle, and syringe. When using this type of setup, the butterfly needle is attached to an extension set and syringe that is filled with the test compound. The tail may be taped to the table and the butterfly needle is then inserted into the vein and placement is verified by aspiration on the syringe. Once confirmed, the butterfly needle may also be taped in place. This prevents the needle from being pulled out of the vein during dosing. This type of setup can be very useful when administering large volumes of test article as a slow bolus over several minutes or when the test compound may be irritating or mildly caustic. Taping the animal's tail in place prevents the animal from pulling the tail out of the fingers of the technician dosing.

If repeated dosing is to be performed, the initial venipunctures should be performed as close to the tip of the tail as possible. During the injection, if the needle comes out of the vein, a bleb will form under the skin. The needle should be repositioned immediately to prevent infiltration of the solution around the vein. Infiltration of an irritating solution can cause necrosis and make future injections difficult or impossible. Injection of 2 mL/100 g body weight can be accomplished without stress to the rat. [Barrow \(1968\)](#) found that injections of volumes over this amount produced respiratory difficulty and pulmonary edema.

Tail vein infusions

Tail vein infusions are convenient because catheter placement can be accomplished without anesthesia. A 23-gauge or smaller needle connected to an extension set is inserted into the tail. The needle and extension set is then secured to the tail with tape. The extension set is

attached to a syringe that is placed on a pump and the test compound can be infused. The tail may be taped to a wooden stick or tongue depressor to further protect the needle from being dislodged. Over-the-needle catheters are also commercially available and offer the advantage that the needle is removed once the catheter is placed in the vein and may help to prevent further penetration of the vein wall and subsequent perivascular dosing ([Rhodes and Patterson, 1979](#)). Advantages that this technique has over permanent indwelling catheters are that the catheter is removed following dosing and will not become occluded and the animal does not have to undergo anesthesia and a surgical procedure to place the catheter. Permanent catheters have a tendency over time to develop a fibrin flap or become clotted, thus losing patency. A major disadvantage is that the animals have to be restrained during the infusion, which may cause stress and alter the results of the study. When using this technique, the duration of the infusion should be limited so that the length of time the animal is restrained is limited.

An alternative technique using the lateral tail vein involves placing a catheter in the vein and wrapping the tail in a similar manner as previously described, and then a lightweight protective cover attached to a tether system is placed around the tail to hold the catheter or needle in place.

Jugular vein

Although this route has been used for bolus injections, it is most widely used as a site for cannulation from indwelling catheters. The indwelling catheter required surgical implantation under anesthesia.

Bolus injection

Although injections can be made by exposing the jugular vein by incision, this method is not acceptable for repeated dosing. The jugular vein can be accessed for test compound administration without exposing the vein. The animal can either be anesthetized or restrained on the back. The head is positioned to either the left or the right for access to the respective jugular vein. A 23-gauge needle fitted to a syringe with the bevel up is inserted in a cephalocaudal direction into the angle made by the neck and shoulder. The needle should enter the vein anterior to the point at which it passes between the pectoralis muscle and the clavicle. When about one-half the length of the needle has penetrated the skin, the bevel should be in the lumen of the vessel. Insertion of the needle through the muscle stabilizes the needle and minimizes bleeding. Caution should be used when using this technique as it is considered to be a "blind stick" into the vessel and damage

to the vessel may occur. Repeated access of the vessel is not recommended.

Infusion

For the purpose of continuous infusion of the test compound over extended periods of time or for repeated short-term infusions, implanted catheters in the jugular vein may be used. For implantation of a jugular catheter, the animal is first anesthetized and placed in a dorsal recumbency and the surgical site is prepared. A midline incision is then made in the neck and a section of the jugular vein is dissected free. Manipulation of the vein should be limited to prevent vasospasm. A cephalic ligature is then tied and the vein elevated. A small incision is then made in the vein and the catheter is passed into the vein and tied in place. The other end of the catheter is then tunneled subcutaneously to between the scapulas where the catheter is exteriorized. The catheter should be filled with an anticoagulant solution such as heparin when not in use. When correctly positioned, the tip of the cannula will be at the junction of both vena cava. If placing catheters into young animals, enough of the catheter should be inserted to allow for growth of the animal. Care should be taken that the catheter is not inserted too far as the tip may be pushed into the right ventricle of the heart. Improper placement of the catheter may lead to administration of the test compound directly into the heart, which can cause complications.

Similarly to the jugular vein, administration of test article via the femoral vein requires an implanted catheter. For implantation of a femoral catheter, the animal is first anesthetized and placed in dorsal recumbency and the surgical site is prepared. A midline incision is then made in the inguinal area and a section of the femoral vein is dissected free. Manipulation of the vein should be limited to prevent vasospasm. A ligature is then tied and the vein elevated. A small incision is then made in the vein and the catheter is passed into the vein and tied in place. The other end of the catheter is then tunneled subcutaneously to between the scapula where the catheter is exteriorized. The catheter should be filled with an anticoagulant solution such as heparin when not in use. When correctly positioned, the tip of the cannula will be positioned in the vena cava. For longer term infusions, the femoral vein catheter may be preferable as patency is easier to maintain and the risk of damage to the heart from the catheter is avoided.

Several commercial vendors offer surgical support services and for an additional fee will implant either jugular or femoral catheters. These vendors will typically have a specific methodology for implant, but will accept requests for modifications such as catheter type and exteriorization site. The typical catheter implanted may be manufactured from polyethylene,

polypropylene, or silastic. In recent years, manufacturers have developed catheters impregnated or ionically bound with heparin. These catheters may help to prolong the life of the catheter.

The useful lifetime for jugular and femoral catheters is quite variable; the lumen of the cannula may eventually become obstructed by a blood clot or fibrous mass. The position of the tip of the catheter is important. Clot formation is less likely to occur if the tip of the catheter is placed in the venous stream rather than in the jugular vein (Popovic and Popovic, 1960). It is recommended for repeated short-term infusions that, when the test article is not being infused, a slow infusion of saline will help to prolong the life of the catheter.

Prior to use, the patency of the catheter should be checked by removing the anticoagulant lock, checking for blood draw back, and then flushing with saline or lactated Ringers solution. Alternatively, patency can be checked by injecting 3–6 mg of pentobarbital solution (0.05–0.10 mL of a 60 mg/mL solution) into the catheter (Weeks, 1972). If the catheter is patent, the rat will lose its righting reflex and become ataxic within 10–15 seconds of injection. The rat will recover in 10–15 minutes.

Rats will destroy the catheter if left between unprotected or in easy reach of the forepaws. By exteriorizing the catheter between the scapula, the rat will not be able to chew on the catheter. For the purpose of continuous infusion, several manufacturers have developed tether systems and catheter sheaths made of metal that prevent the animal from chewing on the catheter. These systems typically consist of a jacket with an attached tether that the catheter passes through. The catheter then attaches to a swivel that prevents the catheter from becoming kinked. The swivel then attaches to a second catheter that can be attached to a syringe or pump for administration of the test compound (Guo and Zhou, 2003). When performing long-term infusion studies, the effects of the catheter and harness should be considered. Infections, septicemia, a variety of visceral lesions, endothelial lesions, and increased platelet consumption have been observed in cannulated animals (Hysell and Abrams, 1967; Meuleman *et al.*, 1980; Vilageliu *et al.*, 1981). Decreased or erratic weight gains and decreased liver and thymus weights have been observed in tethered animals. These changes may be attributed to the stress involved in chronic tethering of the animals.

An alternative to an exteriorized catheter is to attach a subcutaneous port to the catheter that can be accessed via a transcutaneous needle stick. This type of set-up helps to prevent infections that can occur with transcutaneous catheters. One of the pitfalls of this subcutaneous port is that the port may only be accessed a finite number of times. In addition, care has to be taken to ensure the port and catheter are properly flushed of all test compound and blood as clots can easily form.

Administration of small volumes of test compound may be accomplished using a subcutaneously implanted osmotic pump. This type of pump is connected to the catheter after being filled with the test compound and implanted in a subcutaneous pocket. This allows for a continuous administration of small amounts of compound without the need for a jacket and tether system.

Saphenous, lateral marginal, and metatarsal veins

These veins in the leg and foot are easily visualized and can be injected without anesthesia; however, assistance is required. Shaving the area over the saphenous or lateral marginal vein makes visualization easier. During injection it is necessary for one technician to restrain the animal and occlude the vessel to cause it to dilate. Wiping the skin over the vein with 70% alcohol or with gauze soaked in hot water will help to dilate the vessel and increase the possibility of success. The second technician then performs the injection; a 26- to 27-gauge needle should be used.

Dorsal penis vein

When administering test article via the dorsal penis vein, it is preferable to use anesthesia. Lightly anesthetizing the animals with an inhaled anesthetic such as isoflurane or CO₂/O₂ will prevent the animal from struggling and increase the possibility of a successful injection. This procedure requires two technicians to perform the injection. One technician holds the animal by the skin on the back and the feet and tail. The vertebral column is then hyperextended. The second technician then grasps the tip of the penis between the thumb and forefinger, and injects the test solution into the dorsal vein using a 26- to 30-gauge needle.

Sublingual vein

While the method of sublingual vein injection has the disadvantage of requiring anesthesia, it only requires one technician. Ideally, the animal should be anesthetized with an inhaled anesthetic such as isoflurane or CO₂/O₂, but injectable anesthesia may also be used. The animal should be placed in a dorsal recumbency with the head towards the operator. The test compound may be administered by holding the tongue between the thumb and forefinger, and, using a 26- to 30-gauge needle, the vein is entered at a very shallow angle and the injection is performed. After completion of the injection, the bleeding can be stopped using direct pressure. Once the bleeding has stopped, a small cotton-wool pledget should be placed over the vein and the tongue placed back in the mouth. The animal will spit the cotton out upon regaining consciousness.

Intraperitoneal route

Test compounds injected into the peritoneal cavity will be absorbed into the portal circulation and transported to the liver. As a result, the compound will be subjected to the metabolic activity of the liver prior to being circulated to the remainder of the animal. Based upon the level of blood flow and circulatory surface area in the peritoneal cavity, compounds injected intraperitoneally will be absorbed quickly.

Intraperitoneal administration of test compounds in the rat can be performed by one person. The animal should be picked up by the scruff of the neck and back and held firmly in a dorsal recumbency. This position will allow for proper access to the peritoneal cavity. The belly of the animal should be visually divided into quadrants and a needle (<21-gauge) should be inserted anteriorly into one of the lower quadrants just lateral to the midline. Aspiration of the syringe prior to injection will help the investigator to determine if the needle is positioned appropriately.

Intramuscular route

Intramuscular injection of compounds will result in the rapid absorption into general circulation due to the abundant supply of blood vessels. However, the speed of absorption will not be as fast as with an intraperitoneal injection. Acceptable sites in the rat are the quadriceps, the thigh, and the triceps. This procedure can be done with one or two technicians. The selected muscle mass should be stabilized with the thumb and forefinger of one hand while restraining the animal and the needle is guided into the muscle with the other hand. A 21-gauge or smaller needle should be used. The needle should be lightly aspirated to ensure the tip of the needle is not in a blood vessel and then the compound is slowly injected. A slow injection with a minimal volume will help to minimize pain. Approximately 1 mL/kg of solution can be injected per site. If larger volumes are required, multiple injection sites should be used.

Subcutaneous route

Absorption following subcutaneous injection is typically slower than following intramuscular injection. This may be advantageous if a relatively sustained period of absorption is desired. Another advantage of the subcutaneous route versus the intramuscular route is a much larger volume of test compound can be administered. Five to 10 mL can be easily injected with little to no discomfort to the animals. This can be beneficial for test compounds that have limited solubility. Suitable sites for injection are the ventral body, the flank, and shoulders. To perform the injection, the skin is grasped between the thumb and forefinger and raised to make a

tent. The needle (<20 gauge) is inserted through the skin to make the injection. Injection sites can be varied for multiple-dose studies where the solution is a potential irritant.

To minimize the stress of manipulation and to provide a means for continuous infusion, a perforated cannula or catheter can be implanted transcutaneously. The cannula can then be secured using a tether system similar to that used for intravenous infusions. Using this system, the test compound can be infused continuously. Mucha (1980) injected sodium pentobarbital directly into the cannula and showed it was absorbed much more rapidly than following injection. An alternate method of infusion of small volumes over an extended period of time is through the use of an osmotic minipump. The pump is filled with the test compound and implanted subcutaneously under anesthesia. These pumps are commercially available and offer the advantage of a continuous infusion at a constant rate without the animals being encumbered with the infusion apparatus. This method would work only when the solutions are stable at body temperature for the duration of the infusion period. In recent years several types of absorbable microspheres have been developed that can act as carriers for the test compound.

Topical route

The rat has not traditionally been used as a model in skin irritation or sensitization studies. However, the rat has been used in systemic toxicity studies where the skin is used as a portal of entry for whole body exposure or in skin painting studies where the carcinogenic potential is being assessed. In a comparison of absolute absorption rates of several compounds, Bartek *et al.* (1972) found dermal absorption rates in the rat tended to be slightly lower than in the rabbit, and higher than in the monkey, swine, and human.

Exposure in dermal studies is usually to the anterior dorsal portion of the back. The skin should be shaved weekly or 24 hours prior to skin painting. Care should be taken to ensure that the skin is not damaged during shaving, as this can increase the rate of absorption of the test compound. The test area for application should be clearly marked; for repeat dose studies the area of this site is often 10% of the body area. Usually 0.25–1.0 mL of the test solution is applied in skin painting studies. The amount of a cream or ointment applied will vary with the test compound and desired total dose administered. Dosing is typically performed every day.

The actual dose in a dermal toxicity study is determined by the amount of compound absorbed; therefore factors that influence absorption should be considered. Several design features in a topical study may affect absorption: abraded skin will tend to absorb faster than

intact skin; test compound may adhere or build up at the site of exposure and may impede absorption (and test compound may be chemically changed owing to exposure to air or light); the test compound may be licked or scratched from the site; and the test compound may be ingested by the animals.

Several techniques have been developed to avoid removal or ingestion of the test compound. For acute studies, Rice and Ketterer (1977) described a cable-type restrainer attached to a stainless steel plate. Loops just behind the front and just in front of the hind legs hold the animal immobile. Other methods may be used to reduce stress to the animals and allow the animal mobility. One method is the use of an "Elizabethan" collar. This is a 4- to 5-cm wide strip of plastic or other material that fits around the neck of the animals. This prevents the animal from being able to turn its head to gain access to its back. Consideration should be given to the use of this method as the collar also may prevent the animal from being able to properly eat from a feed jar. A second method is to wrap the animal with gauze and then with plastic wrap. Care should be taken when wrapping the animal to avoid excessive tightness. This type of covered exposure may affect the absorption of the test compound. Other types of harness, collars, and acrylic chambers may be used and the appropriate technique should be chosen based upon the intended length of exposure and the efficacy with which the technique can be performed by the technicians.

Rectal route

The rectal route is not a routinely used method of administration in toxicology. However, administration by this route is sometimes required to support drugs given rectally by suppository. For dosing, the animal is held by the base of the tail and a stainless steel, ball-tipped gavage needle (5 cm) or vinyl tube (6 cm) attached to a syringe is inserted into the rectum. Care must be taken not to damage the rectum when inserting the needle. The syringe should be held lightly: the weight of the needle and syringe propel the needle. The animal can either be awake or anesthetized. If animals are awake, excretion of the unabsorbed test compound may occur. Methods to control excretion have included ligation of the rectum (Nishihata *et al.*, 1984) or various types of septums which are tied or glued in place (DeBoer *et al.*, 1980; Iwamoto and Watanabe, 1985). Anesthetized animals can be placed on an inclined board to retard expulsion.

An important factor in rectally dosing the rat is that the depth of deposition of the test compound will affect the rate of absorption and should be standardized. Drugs subject to extensive first-pass metabolism, such as propranolol and lidocaine, have been found to be

much more bioavailable when injected close to the anus rather than in the upper areas of the rectum (DeBoer *et al.*, 1980; Iwamoto and Watanabe, 1985). The reason for this difference in bioavailability appears to be that the venous return in the upper rectal area is through the upper rectal vein that feeds back into the portal circulation and then into the liver. The venous return in the lower rectum is through the lower hemorrhoidal veins and is not connected to the portal system, but goes directly to the inferior vena cava (Iwamoto and Watanabe, 1985).

Intranasal route

With the increasing number of drugs being delivered nasally, methods have been developed to support this route. For administration in unanesthetized animals, the appropriate volume of test material is drawn into the tip of a pipette or other appropriate dosing implement. The tip of the dosing implement is placed directly over but not into the nostril to be dosed and the test material is instilled into the nostril. The animal will aspirate the test material into the nasal passage. This can be repeated for the opposite nostril, or the opposite nostril can be used as a control treatment. In the event that the animal sneezes or the test compound is otherwise expelled, the nostril should be treated a second time.

Inhalation route

Owing to the complexities and equipment involved in generating, maintaining, and measuring appropriate atmospheres, the inhalation study is one of the most technically difficult to perform. This section is not written to provide a complete discussion of the skills necessary to perform an inhalation study, but will deal with the general considerations about the three major steps in exposing rats by the inhalation route; generation of the test atmosphere, exposure of the test animals, and measurement and characterization of the test atmosphere.

Intratracheal administration

An alternative to inhalation administration is to instill the test compound into the trachea of the animal. Techniques have been developed to instill the test compound into the trachea safely and repeatedly. To perform the procedure, the animal must first be anesthetized, preferably with a gas anesthesia such as isoflurane. Once it is anesthetized, a speculum is inserted into the mouth and passed into the trachea. A syringe and needle with a 5 cm piece of tubing is used to instill the test compound. The tubing is then passed over the speculum and the test compound is administered. Volumes should be limited to 2 mL or less of test compound.

End point measurement techniques

Observations and physical examinations

Rats are routinely monitored during toxicology studies as an assessment of their general health and to define the effects of the test article. In acute and subchronic studies, animals may be observed frequently in an effort to define short-term pharmacologic changes induced by the test compound, which may become apparent at peak blood levels. Specifically in acute toxicology studies clinical observations will help to establish a maximum tolerated dose. In chronic studies, these observations are critical in tracking tumor development and for determining animals in extremis, which should be euthanized for humane reasons and to prevent autolysis and tissue loss. Arnold *et al.* (1977) provide a useful description of a clinical assessment program for chronic studies.

Daily observations are performed first thing in the morning and last thing before leaving in the afternoon to assess the health of the animals and identify animals that may be *in extremis*. In this observation, behavioral status, respiratory signs, skin, eyes, and excretory products are noted. Care should be taken to disturb the animal as little as possible, as this may induce stress and affect the animal's behavior. The animal should be picked up and examined more closely if abnormalities are detected. Special attention should be paid to the amount of feces present, since a decrease in fecal output may be the first signs of a watering system malfunction. In acute studies or where pharmacological effects are expected, animals may be examined continuously or at peak plasma levels.

A more thorough physical examination should be done weekly. Each animal is taken from its cage and placed on an examination table where respiration, behavior, general appearance, and locomotion are observed. The technician should then pick up the animal, examine its body orifices, skin, and coat, and perform a palpation of the trunk and limbs to check for tumors. The detection and tracking of the size and fate of masses (potential tumors) is essential for carcinogenicity studies.

Animals that have experienced severe weight loss over the previous week, a progressive decline in weight over several weeks, or other severe clinical signs should be observed more frequently and marked for possible euthanasia. Body and feeder weights can be measured as part of the physical examination or as a separate function. Performance of these operations as a part of physical examinations will minimize animal handling and potential stress. If the operations are combined, it is important that the physical examinations be done completely and are not rushed.

Neurobehavioral examination

Neurobehavioral examinations are included in toxicology studies to assess the behavioral and neurological effects of test compounds. These examinations, which may be done as part of acute or repeat dose toxicity studies or a study specifically designed to assess just the neurobehavioral effects, typically involve screens consisting of an abbreviated functional observational battery and some measure of locomotor activity. The Environmental Protection Agency (EPA) has written guidelines on the design and conduct of these studies (OECD, 1995a,b). To meet EPA guidelines, the screen is performed prior to the start of treatment, then periodically during the course of treatment. It may be performed as a separate study, on satellite groups of animals in conjunction with the main study, or on animals in the main portion of the toxicology study. Recently, the ICH has adapted guidelines requiring neurobehavioral assessment of all new pharmaceuticals prior to initiation of Phase I clinical trials. To meet ICH guidelines, a more formal functional observational battery should be conducted and may be conducted on main study animals or as a separate study. Where initial screens indicate the possibility of a test compound-related change, a more specialized series of tests may be performed to assess the nature of the effect and the extent of the central nervous system involvement. These secondary tests evaluate motor and sensory function as well as cognitive ability. Examples of secondary tests include sensory-evoked potential experiments and schedule-controlled behavior studies. Descriptions of these secondary tests can be found in [Annau \(1986\)](#).

Functional observational battery

The typical observation battery (FOB) includes observation of home-cage and open-field activity as well as measurements of reflexive, physiological, and neuromuscular function ([Gad and Chengelis, 1998](#); [Moser et al., 1988](#); [Moser and Ross, 1996](#)) as outlined in [Table 2.15](#). Observations and measurements that have become standard for an FOB evaluation are presented in [Table 2.16](#). The order of measurement should be consistent, progressing from the least interactive to the most interactive measurements. Home-cage observations are made first. Assessments of posture, clonic movements, tonic movements, and palpebral closure (if the animal's eyes can be seen) are taken prior to removing the animal from its cage. However, it may be necessary to pull the cage from the cage bank or remove the cage cover in order to see the animal's eyes. The animals are then transferred to the open field. During the transfer, certain physical observations are made. The technician removes the animal from the cage. The

technician holds the animal and notes increased or decreased body tone as well as such observations as bite marks, soiled fur appearance, missing toenails, emaciation (shallow stomach, prominent spinal vertebrae), or death. In addition, observations of lacrimation, palpebral closure, piloerection, exophthalmos, and salivation are also made. The animal is then placed in the open field apparatus for a set period of time. Measurements of rearing, urination, and defecation are made immediately at the end of the assessment period. Assessment of clonic movements, tonic movements, gait, mobility score, arousal, respiration, stereotypic behavior, and bizarre behavior may be made immediately after the time period has ended or the technician may continue to observe the animal for a longer period of time to allow for more accurate assessment. When the open field assessment has been completed, the animal is removed from the open field apparatus for the

TABLE 2.15 Functional observational battery in rats

	Number of Animals	
	Males	Females
Control	10	10
Low Dose	10	10
Mid Dose	10	10
High Dose	10	10

DOSING: The test article will be administered by the required route.

FUNCTIONAL OBSERVATIONAL BATTERY: FOB evaluations will be conducted prior to dosing, at the estimated time of peak effect, and 24 hours postdose. The evaluations are as follows:

Home Cage Observations:

Assessments of posture, clonic movements, tonic movements, and palpebral closure (if the animal's eyes can be seen) are taken prior to removing the animal from its cage.

Handling Observations:

Observations of ease of removal, handling reactivity, lacrimation, palpebral closure, piloerection, and salivation are made upon removal of the animal from the home cage.

Open Field Observations:

The animal is placed in the center of an open-field testing box (measuring 20" × 20" × 8"). Clean absorbent paper may be used to cover the bottom of the box if required by protocol. Using a stopwatch, the animal's stay in the box is timed for 3 minutes. Measurements of rearing, urination, and defecation are made immediately at the end of the 3 minutes. Assessment of clonic movements, tonic movements, gait, mobility score, arousal, vocalization, respiration, stereotypic behavior, and bizarre behavior may be made immediately after the 3 minutes have ended or the technician may continue to observe the animal for a longer period of time to allow for more accurate assessment.

Sensorimotor Observations:

The approach response, touch response, click response, and tail-pinch response (stimulus reactivity tests) are performed while the animal is in the open field apparatus, after the 3-minute time period is over and all other measurements have been recorded. The animal is removed from the open field apparatus for the pupil response, righting reflex, thermal response, hindlimb splay, and grip strength measurements.

Physiological Evaluations:

The animal's body weight and rectal temperature are measured and recorded.

CLINICAL EXAMINATION: Following each FOB assessment (additional observations will be conducted prior to dosing for locomotor activity animals, if requested).

TABLE 2.16 Functional observation battery

Home Cage	Open Field	Manipulative	Physiological and Neuromuscular
Posture	Rearing	Ease of removal from cage	Body weight
Clonic movements	Urination	Handling reactivity	Body temperature
Tonic movements	Defecation	Lacrimation	Hindlimb extensor strength
Palpebral closure	Clonic movements	Palpebral closure	Grip strength
	Tonic movements	Piloerection	Hindlimb splay
	Gait	Exophthalmos	
	Mobility	Salivation	
	Ataxia	Approach response	
	Arousal	Touch response	
	Vocalizations	Click response	
	Respiration	Tail pinch response	
	Stereotypy	Pupil response	
	Bizarre behavior	Eye blink response	
		Forelimb extension	
		Hindlimb extension	
		Righting reflex	
		Thermal response	

Source: From Gad (1982), Moser *et al.* (1988), and Haggerty (1989).

approach response, touch response, click response, tail pinch response, pupil response, righting reflex, thermal response, hindlimb splay, and grip strength measurements. After completion of the manipulative assessments, the physiological evaluations are completed.

Procedures used during the performance of an FOB must be standardized because some observations made have a subjective component. If at all possible, a single observer should be used throughout a single study. If not possible, a single observer should at least conduct all assessments of an animal. In addition, technicians should be blinded to the treatment conditions for each animal.

Locomotor activity

Methods used for recording motor activity include direct observation and automated techniques such as photocell devices and mechanical measurements (Macphail *et al.*, 1989). In direct observations, the observer can make quantitative measurements of the frequency, duration, or sequencing of various motor components of behavior or qualitative records on the presence or absence of certain components of activity. Photocell devices record the number of times an animal interrupts a beam in specially designed chambers. In mechanical chambers, the animal's movements result in a vertical or horizontal displacement of the chamber; records are kept of the chamber's movements. There are advantages and disadvantages of each technique. In direct observation, a record can be made of behavior, such as convulsions, that may not be observed when using the photocell or mechanical methods. A disadvantage of the direct observation method is that the animal

may be influenced by the presence of the observer. Advantages to the photocell and mechanical methods are that the data is captured electronically, the observer does not have to be present, and the computer system can graphically present the data in the form of lines crossed or a map of the activity.

To make activity determinations, an animal or group of animals is put into an observation or recording chamber and activity is recorded for a specific period of time. Since activity will normally decline over the course of the session, the length of the observation period is important. The EPA guideline specifies that activity should approach asymptotic levels by the last 20% of the session. Haggerty (1989) used a 15-minute recording session, accumulating data over three 5-minute intervals. Since a large number of environmental conditions can affect motor activity, e.g. sound level, cage design, lighting, temperature, and humidity, or odors, it is important to minimize variations in the test environment.

Cardiovascular parameters

Examinations of the cardiovascular system may be scheduled into toxicology studies or performed when the cardiovascular system is a suspected target of the test compound. The ICH has adapted guidelines requiring cardiovascular safety assessment of all new pharmaceuticals prior to initiation of Phase I clinical trials. This guideline recommends that this assessment be conducted in a nonrodent species, but cardiovascular assessment can be performed for screening purposes or as additional support data.

Electrocardiography

Although the dog has traditionally been the species of choice in toxicology studies of effects of ECG, research with the rat has progressed and increased over the years. [Detweiler \(1981\)](#); see also [Detweiler et al., 1981](#) provides an excellent review of the use of electrocardiography in toxicology studies in the rat. This section will present a general discussion on the aspects of the recording methods and interpretation of the ECG in the rat.

Recording methods

Restraint: One of the disadvantages of traditional methods of studying ECGs in rats is that it is difficult to keep the animals still while recording. It is important that the animal remain in a constant position during the procedure using skin leads to avoid muscular artifact in the ECG. Various forms of restraint have been tested, each of which requires acclimation to the procedure prior to evaluation. These methods included restraining the rat in a supine position using rubber gloves ([Hundley et al., 1945](#)), pinning the animal to a board, boards with clamps, and plastic tubes with slits on either side which allow for placement of the electrodes ([Zbinden et al., 1980](#)). Various forms of anesthesia have also been evaluated.

No conclusion has been reached about the best method of restraint. The basic concerns are that manual methods and physical restraint require acclimation to allow the animal to become accustomed to the procedure and tracings can be reasonably free of muscular artifact. Also, varying pressures of clamps or handling during the restraint may affect the results. The use of anesthesia has been shown to produce changes in the ECG and the possibility of drug interactions between the anesthetic and the test compound may occur.

Position: The most common positions are the prone or ventral recumbency position when animals are awake and the supine or dorsal recumbency position when anesthetized. [Beinfeld and Lehr \(1956\)](#) compared the positions and concluded that the prone position produced an increased R wave and it avoided unfavorable cardiac rotation and an undesirable variation in the projection of the special QRS loop.

Tethered: [Robineau \(1988\)](#) developed an electrode system which can be implanted subcutaneously a few days prior to recording. The device has disconnected that is exteriorized between the scapula. The advantage of this system is that a cable can be connected to the plug and the ECG can be taken in unrestrained rats. The disadvantage of this method is that it should only be used for short-term studies as the implant provides a source for infection in the animal.

Telemetry: Recently, techniques have been developed for monitoring cardiovascular parameters via telemetry

in the rat ([Kuwahara et al., 1994](#); [Kramer et al., 1995](#); [Ichimaru and Kuwaki, 1998](#)). Totally implantable battery operated systems have been developed that can monitor several physiological parameters including ECG. The implants are available with ECG leads that can be positioned in the Lead II configuration for monitoring ECG continuously for extended periods of time in a freely moving animal. As the leads are attached subcutaneously to the musculature, the signal is of a higher quality than that gained using skin leads. Also, the animal is not affected by the observer, as it can be monitored remotely. One of the disadvantages is that these implants are only designed to monitor a single lead.

Leads: Most investigators use Einthoven's bipolar limb lead system, lead I (right and left foreleg), lead II (right foreleg and left hindleg), and lead III (left foreleg and left hindleg), with and without the augmented unipolar limb leads aVR (right foreleg), aVL (left foreleg), and aVF (one of the hindlegs). Since the foreleg position can alter the scalar ECG wave amplitudes, investigators must standardize these positions during recording. When implanting telemetry leads, the leads are placed on the right clavical and the most caudal rib on the left side in a modified lead II configuration.

In the past, various types of leads have been used to connect the ECG wires. These included the use of hypodermic needles inserted under the skin; small gauge insulated copper wires wrapped around the shaved distal portion of the limbs; or alligator clips. In addition, platinum tipped pin electrodes are commercially available that provide a good quality signal with limited discomfort to the animal.

ECG waveform

The major points to notice about the rat ECG are that the conventional waves of the mammalian ECG (P, QRS, and T) are all identifiable in the rat ECG, there is no isoelectric line during the electrocardiographic complex, and there is no ST segment. The duration of the standard intervals evaluated in the ECG of the rat are as follows: P, 10–20; PR Interval 35–50; QRS complex 12–25; and QT interval 38–80 ms ([Detweiler, 1981](#)). The duration of the intervals is related to the heart rate: as the heart rate increases, the intervals shorten and as the heart rate slows, the intervals prolong. These intervals can also be affected by administration of test compounds and are the basis for the requirement to assess the effects of a test compound on cardiac function. Specifically, prolongation of the QT interval has been correlated with a phenomenon called *Torsade de pointes* which can lead to sudden cardiac death. Several classes of compounds, such as antihistamine and Ca²⁺ channel blockers, have been shown to prolong the QT interval ([Gras and Llenas, 1999](#)). Each wave of the ECG represents either a depolarization or repolarization of the

atria and ventricles of the heart. For example, the P wave is an electrical representation of the depolarization of the right atria and the T wave is an electrical representation of the repolarization of the left ventricle. Spear (1982) provides a more in-depth discussion of the waveforms and the electrophysiology of the heart. Several computerized systems have been developed that are capable of recognizing the independent waveforms and measuring the intervals. Caution should be used when using these systems as they are typically programmed to recognize a normal ECG and the presence of arrhythmias may be missed or the interval may not be measured correctly.

Heart rate

The heart rate can be calculated from standard limb lead ECGs by measuring the distance between the two peaks of the R wave. This distance is then divided by the chart speed (i.e. 50 mm/sec) to calculate the RR interval in seconds. This is then divided into 60 seconds to calculate the heart rate in beats per minute. Using the formulas below, the heart rate of an RR interval of 10 mm measured on a chart printed at 50 mm/sec = 300 beats per minute.

$$\begin{aligned} & \text{RR Interval (mm/beat)/Chart Speed (mm/sec)} \\ & = \text{RR interval (sec/beat)} \\ & (60 \text{ sec/min})/\text{RR interval (sec/beat)} \\ & = \text{Heart Rate (Beats/min)} \end{aligned}$$

Many environmental things can affect the heart rate of the rat, such as excessive manipulation, technicians the animal is not familiar with, new environments, etc. Therefore, when evaluating ECG using a restrained method, it is important to acclimate the animal to the test procedures prior to starting. Detweiler (1981), in a review of the literature, found that published heart rates for rats varied between 250 and 750 beats/minute. Awake, restrained adult rats had heart rates from 330 to 600 beats/minute and well acclimated restrained adult animals had heart rates from 250 to 350 beats/minute. Heart rate in unrestrained telemeterized animals has been reported to be between 225 and 350 beats/minute (Zbinden, 1981; Guiol *et al.*, 1992).

Blood pressure

During the conduct of a toxicology study it may be necessary to monitor blood pressure. This can be done using either indirect or direct methods. Caution should be used when using indirect methods as the values obtained may be variable. The direct method involves the implantation of an arterial cannula for measurement of blood pressure. This method is more reliable, but has a limited period of time during which the cannula may remain patent. It is recommended that for definitive

assessment of the hemodynamic effects of a test compound, a nonrodent species such as a dog or a nonhuman primate be used.

Indirect measurement

Indirect methods of blood pressure measurement detect systolic blood pressures by the occlusion of arterial inflow of blood and the subsequent detection of the pressure at which the first arterial pulsation occurs. The two places where indirect measurements can be made on the rat are the tail and the hindpaw.

Tail Cuff Method: The tail cuff method monitors pressures in the ventral caudal artery. In this method, the animal is put into a restrainer that allows for free access to the tail. An inflatable cuff is then placed around the base of the tail and the pressure is increased until flow stops, and the pressure is then slowly released until flow resumes. The cuff pressure at the time when flow resumes is the systolic blood pressure. Various methods have been used to determine when this occurs.

Because the caudal pulse is rather weak, preheating of the animals in boxes at temperatures of 30–42°C for periods up to 10 minutes may be necessary, to dilate the caudal artery. This technique should be used with caution, as previously discussed changes in body temperature can have widespread effects on the animal and may produce unexpected test compound effects.

The placement and width of the tail cuff is important. There is a gradient in pressure along the caudal artery which amounts to 4.5 mmHg/cm. For this reason, the cuff should be placed close to the base of the tail and this should be standardized. If multiple readings are done over time, marking the placement of the cuff with an indelible marker will help to standardize placement. In addition, variation in the width of the rubber tubing can be a source of error. Bunag (1973) found that the most accurate readings were given by a 15-mm cuff; shorter cuffs gave falsely elevated readings and longer cuffs gave low readings.

Hindpaw Method: Measurement of blood pressure in the hindpaw does not measure the pressure in a specific vessel. In this method, the animal is placed in a restrainer, as in the tail cuff method. A pressure cuff is placed around the ankle to occlude blood flow and blood pressure is measured as the cuff pressure is released and blood flow returns. As in the tail cuff method, several techniques have been used to determine the return of blood flow. These include visual observation (Griffith, 1934), photoelectric cell (Kersten *et al.*, 1947), and oximeter (Korol and McShane, 1963). The advantage of the photoelectric and oximeter methods is that they do not require preheating to dilate the vessels of the hindpaw. The oximeter method measures mean arterial pressure rather than systolic pressure.

None of the indirect measurements is able to evaluate the complete hemodynamic cycle. They evaluate only systolic or mean arterial pressure, but not systolic, diastolic, and mean arterial pressure together.

Direct measurement

The direct measurement techniques involve the cannulation of an artery with the blood pressure being determined with a manometer or transducer connected to the free end of the cannula. This is true for types of direct measurement including telemetry. Surgery and cannula placement utilize similar techniques as those previously described for placement of a venous catheter. In short, the artery is isolated and a small incision is made in the femoral artery. The catheter is then inserted into the hole and passed into the abdominal aorta. The carotid artery may also be used for this procedure, but care should be taken not to insert too far as the catheter may be passed into the left ventricle of the heart. If this occurs, the blood pressure waveform will change in appearance. The left ventricular waveform shows a similar systolic pressure as does an arterial blood pressure waveform. But the diastolic pressure is much different. If the catheter has been placed in the left ventricle, the diastolic pressure will be 0 mmHg or slightly negative. If this occurs the catheter should be backed out into the aortic arch.

Where chronic use is desired, cannulas are typically run subcutaneously and exteriorized between the scapulas or at the back of the head. A carotid artery catheter can be expected to remain patent for 3–5 weeks (Ross, 1977; Andrews *et al.*, 1978), while an abdominal aorta catheter may remain patent for several months. Care should be taken when using arterial catheters for long periods of time as fibrin deposits can build up on the catheter or clots may form in the catheters. The risk exists that these deposits or clots could be expelled during the flushing of the catheter or during normal movement of the animal. These clots or deposit, once free, may occlude other vessels downstream and in the case of a carotid catheter may cause a stroke to occur.

Blood collection techniques

Blood samples are routinely collected in safety studies to determine (1) direct test compound effects on the blood or bone marrow, (2) effects on other organs as indicated by the contents of the blood, for instance, leakage enzymes such as aspartate aminotransferase, and (3) blood levels of the test compound or its metabolites. A variety of techniques have been described for the collection of blood from the rat. The choice of a specific technique may depend upon factors such as (1) the volume to be collected, (2) if the animal is to survive the procedure, (3) the frequency with which samples will need to be collected, (4) whether anesthetics can be used, (5) likelihood

of the animal surviving the procedure, and (6) the impact of organ damage resulting from the procedure. An adult rat has a blood volume of about 50 mL/kg; approximately 10% of the total blood volume can be collected from a rat in a single draw without adversely affecting hematology parameters. For longer term studies, 10 mL per kg per 2 weeks is a reliable guideline for volume of blood drawn. If these volumes are exceeded, hematocrit and red cell mass may be reduced upon evaluation (Diehl *et al.*, 2001).

Technique, anesthetic used for blood draws, and treatment of the animals (i.e. fasted or not fasted) should be standardized throughout a study if repeated samples are being taken. The technique, anesthetic, and handling of the animal may produce effects on the hematology or clinical chemistry parameters to be evaluated.

Retro-orbital plexus

The retro-orbital plexus is a commonly used site for periodic sampling during the course of a study. This method has been shown to be a reliable one for the repeated collection of blood samples. Collection from this site should always be conducted under anesthesia to reduce pain and stress to the animal. Light anesthesia with a mixture of carbon dioxide and oxygen will minimize struggling of the animal and will help to ensure a quick collection with little injury to the animal.

Blood is collected using a microcapillary tube or the fine end of a Pasteur pipette. The tube is inserted into the orbit of the eye at an anterior angle formed by the lids and the nictitating membrane. A short thrust past the eyeball will make the tube enter the slightly resistant horny membrane of the sinus. The tube may be rotated slightly as it is inserted. Once the sinus has been punctured, blood will fill the tube. Once the tube is filled the blood may be allowed to drip out of the end of the tube into an appropriate collection tube. If the flow stops, the tube may be pulled out or advanced slightly to reestablish flow.

In the hands of an experienced technician, there is minimal risk to the animal. Studies have shown that repeated collection of blood from the retro-orbital sinus can produce histological and behavioral changes that may require an animal to be removed from study (McGee and Maronpot, 1979). Several serious side effects of this collection method have been documented. These include retro-orbital hemorrhage, corneal ulceration, keratitis, pannus formation, damage to the optic nerve, and fracture of the orbital bones. When designing the study, the method of blood collection should be taken into consideration. If an important endpoint of the study is ophthalmological examinations, this method should not be used. In addition, anesthetization of the animals may affect other study endpoints.

Tail

A tail vein bleed offers a visible target and is of minimal risk to the animal. Blood will flow faster if the tail has been warmed causing vasodilation. This may be accomplished by dipping the tail in warm water (40–45°C), placing the animal in a warming cabinet for 5 to 10 minutes, or by warming the tail with a heat lamp. This method does not require that the animal be anesthetized, but it should be restrained such that the tail is held immobile. Several methods may be employed to collect blood from the tail. These include clipping the end of the tail off, and vein puncture or artery puncture.

Tail clip

For this method, the animal may be lightly anesthetized or placed in a restraint tube. To collect blood, 2–3 mm of the distal part of the tail is amputated with sharp scissors or a scalpel blade. Three to 4 mL of blood can be collected in 20–30 seconds from a 200–250 g rat. When collection has been completed, the cut surface may be cauterized with a hot spatula or glass rod. This method produces a reliable volume of blood, but should not be used for repeated sampling over extended periods of time.

Venipuncture

Animals should be restrained in a holder that allows complete access to the tail. The tail should then be cleaned and pressure may be applied to the base of the tail causing the vein to dilate. The vein should be punctured using a 21-gauge needle and the blood can either be slowly withdrawn into a syringe through the needle or the needle may be removed and the tail allowed to bleed freely. The use of a butterfly needle may help facilitate the collection of blood as it is less likely to be dislodged if the tail moves. Collection of blood from the tail vein typically yields samples of between 0.5 and 1.0 mL. Upon completion of sample collection the needle should be withdrawn and pressure applied to the tail to stop the bleeding (Frank *et al.*, 1991).

Arterial puncture

This method is conducted in a similar manner to a venipuncture in that the animal is placed in an appropriate restraint tube and the tail is then cleaned. A 21-gauge needle is then inserted into the artery in the midventral surface of the tail close to the distal end. The blood may be then withdrawn into a syringe or allowed to flow freely into a collection tube. The animal's blood pressure will insure the blood continues to flow until pressure is applied to the wound.

Cardiac puncture

This method should always be performed under general anesthesia and should not be used as a survival

collection technique. This technique offers a rapid method for collection of a large volume of blood from the rat. It is possible to collect between 5 and 7 mL from a 300–350 gram rat using this technique and it is possible to exsanguinate the animal using this method.

To collect blood using this method the animal should be anesthetized with a combination of carbon dioxide and oxygen or carbon dioxide alone. The animal is then placed in a dorsal recumbency and the heart is located. The heart may be located by placing the index finger over the 4th and 5th left ribs and the thumb on the right side of the thorax. The collection needle (25–26 gauge, 1–2 cm long) should be inserted at a 45-degree angle into the heart. Once the needle is introduced, the syringe should be aspirated slightly to produce a vacuum. The needle is then advanced until blood is obtained.

Abdominal aorta and vena cava

Collection from the abdominal aorta offers a convenient way to exsanguinate an animal and obtain a maximal amount of blood. To perform the procedure, the animal is anesthetized and the aorta exposed by dissection. A section of the aorta distal to the diaphragm is then exposed and the proximal end is clamped. The aorta is then cut and the distal end placed in a collection tube and the clamp is released. This method will allow for collection of a maximal amount of blood in a short period of time. This method should only be used as a terminal procedure. Alternatively, the aorta may be accessed using a needle or a butterfly needle and samples may then be collected into a syringe.

Winsett *et al.* (1985) described a method of repeated sampling from the vena cava of conscious rats. An assistant holds the animal while the operator grasps the animal just below the last rib. The needle is inserted 1 cm to the right of the spinous process of the first lumbar vertebra at a 45-degree angle until the needle touches the bone. The needle is then slightly withdrawn and then advanced at a slightly shallower angle to miss the bone and access the vena cava. The maneuver of first identifying the bone is essential to the procedure. This procedure may be used for repeated collection, but care should be taken to ensure the animal does not struggle during the collection. If the rat struggles the needle may lacerate the vena cava, causing the death of the animal.

Jugular vein

The jugular vein provides a means for chronic blood sampling that is of low risk to the animal's health. This method does not require the use of anesthesia and can be accomplished through the proper restraint and positioning of the rat.

The unanesthetized method requires two technicians to perform the collection. The animal is placed on a restraint board in a dorsal recumbency. The forelimbs are tied down to the board and one technician holds the hindlimbs of the animal. The second technician grasps the animal's head and turns it down and away from the desired collection site, right or left jugular. The needle (21-gauge) is then inserted into the middle of the triangle formed by the neck, shoulder, and clavicle, parallel to the body. As the needle is inserted the syringe is aspirated until blood is observed in the syringe. The collection is then completed and the site is held off for approximately 30 seconds. It is important that the technician holding the hindlimbs of the animal continually observe the respiratory rate of the animal. If the head is turned too far or is held in the wrong position for too long, the animal may go into respiratory distress. This method may be used for repeated collections with high levels of success.

Alternatively, the animal may be anesthetized and the jugular vein may be surgically exposed. The animal should be prepared using standard aseptic technique and the ventral neck should be shaved. The jugular vein may be exposed by incision of the skin and dissection of the subcutaneous tissues. A needle (20-gauge) may then be inserted into the vein through the pectoral muscle and directing the needle towards the head. Inserting the needle through the muscle will help to stabilize the needle during the collection. Once the needle is in place, the blood should flow freely with little to no aspiration. Care should be taken when aspirating the syringe as too much pressure will cause the vein to collapse. Once the collection is complete, the incision can be closed with a wound clip. This method may be used as an alternative if the animal does not require serial bleeds or as a method to replace a terminal bleed. If the procedure is used for a terminal bleed, strict aseptic technique is not necessary.

Proximal saphenous and metatarsal vein

A small amount of blood (0.1–0.2 mL) can be collected from an animal at minimal risk to their health by utilizing the proximal saphenous and metatarsal veins. No anesthesia is required.

Proximal saphenous vein

The inner aspect of the thigh of the hind limb should be shaved free of hair. While one technician holds the animal and compresses the inguinal area to dilate the vein, a second technician creates a longitudinal nick in the vein with a 20-gauge needle or a hematocrit lancet. The blood can then be collected into heparinized capillary tubes. This method works well for repeated sampling of small amounts of blood but would not be appropriate

for blood volumes required for evaluation of clinical pathology parameters.

Metatarsal vein

This procedure can be conducted with or without an assistant. The animal is restrained and a nick is made in the vessel with a needle. The blood can then be collected into a capillary tube or through the needle into a syringe.

Sublingual vein

When using the sublingual vein for blood sampling, the animal should be anesthetized and then cradled in the palm of the hand. By holding the animal's head between the thumb and index fingers, the head can be stretched back and the skin of the face pulled backwards. This will force the mouth open and the tongue against the palate. The right or left vein should be cut with iris scissors and the animal held such that blood drips into the collection tube. The bleeding can then be stopped by applying pressure with a gauze pad or cotton tipped applicator.

Decapitation

Decapitation should only be performed by trained technicians with the appropriate equipment. There are several commercially available small animal guillotines that should be used to perform this technique. This technique is appropriate when a maximal blood volume is desired and contamination of the sample is not considered to be an issue. To perform this method, the head is first removed and the animal is held over the collection vessel and arterial and venous blood is allowed to drain from the body.

Cannulation

While the blood collection methods described above will provide sufficient volumes and quality of sample for the majority of toxicology studies, specific protocols may require blood to be sampled from rats that have been subjected to a minimum of handling or from specific sites within the body of the animal. Cannulation of a specific artery or vein will typically meet this requirement, though in rodents, the usable life span of the cannula is limited by the length of time the cannula remains patent. Yoburn *et al.* (1984) compared jugular, carotid, and femoral cannulas for long-term sampling of blood. They found the femoral artery cannula was preferable in terms of patency and postsurgical weight loss. Collection from a cannula is the same regardless of implant site. The cannula is typically exteriorized between the animal's scapula and a stylet is inserted into the end of the cannula. For the purpose of

collection, the stylet is removed and a needle is inserted into the cannula with a syringe attached. The heparin lock is then drawn out of the cannula until blood is observed in the syringe. The syringe is then removed from the needle and a new syringe is used to collect the sample. The collection syringe is then removed from the needle and replaced with a syringe filled with the desired solution for locking the cannula, typically a heparin dextrose solution. Once the cannula is flushed and locked with the heparin solution, the stylet is replaced. Using the appropriate technique for flushing and locking the cannula is the key to maintaining its patency.

Jugular vein

The cannulation procedure is the same as previously described for infusion techniques. The cannula has been found to remain patent for variable periods of time and the length of time it remains patent is directly related to the skill of the technician collecting the samples. It is important to remember that a cannula placed in the venous system can easily develop clots in the cannula if not flushed and locked properly. Various methods of anchoring the cannula to the rat's back or head have been developed for ease of sampling or as a connection point for continuous infusion. Each of these methods has been developed so that the exteriorized cannula is positioned such that the animal cannot gain access to it.

Inferior vena cava

The inferior vena cava appears to provide a site for long-lived cannulas. The cannula can be surgically placed either directly into the vena cava through an abdominal surgery or can be advanced into the vena cava from the femoral vein. Either way, the cannula has been shown to remain patent for months (Kaufman, 1980). In either implantation, the cannula is then tunneled subcutaneously and exteriorized between the scapulas in the same manner as the jugular vein cannula.

Abdominal aorta

The most common method for placing a cannula in the abdominal aorta is via the femoral artery. The cannula is inserted into the femoral artery and advanced to the level of just above the kidneys. The opposite end of the cannula is then tunneled to and exteriorized between the scapulas. This is a minor surgical procedure for the animal that the animal recovers from easily. Similar methods are employed when collecting blood from the arterial catheter as those used for the venous catheter. Caution should be taken when removing the stylet from the cannula, as a cannula in the arterial system is under pressure. If not properly prepared, the animal could quickly lose a large amount of blood. It is recommended that the technician place a clamp on the cannula prior

to removal of the stylet. Once the blood collection is complete, the cannula should be thoroughly flushed with saline prior to locking the cannula. The tip of an arterial cannula is positioned such that it is against the flow of blood. Due to this, clots and fibrin deposits develop easily on the end of the cannula, limiting the usable life.

Subcutaneous ports

An exteriorized cannula is a source of contamination and infection, and is subject to destruction by the animal or other animals if group housed. Several types of subcutaneous ports have been developed for implantation along with the cannula. These ports are designed such that test materials may be injected or infused through them or blood may be collected from the port. In most cases, the port is implanted subcutaneously on the dorsal side of the animal. As with exteriorized cannula, the skill of the technician accessing, collecting samples, flushing, and locking the port directly affects the usable life of the port.

Urine collection

Urine is generally collected in toxicology studies to assess kidney function. The most common method used for urine collection is a stainless steel commercially available cage. The cage is designed such that the urine and feces are separated by a cone-shaped device. The urine drains off the collecting walls into a tube and the feces fall into an inverted cone. Food and water are made available in such a way that the urine will not be contaminated. While this type of cage produces urine of acceptable quality for normal urinalysis, the sample may be contaminated with hair or feces. Other methods for urine collection in the rat include cystocentesis, which involves insertion of a needle stick into the bladder, and cannulation of the bladder.

Necropsy

The necropsy is the link between antemortem findings and histological observation. It is an essential portion of the toxicology study and since a necropsy will involve the processing of a large number of animals, it is important that the procedure be well planned (Black, 1986). At a pre-necropsy meeting involving the pathologist, prosectors, and the study director, necropsy responsibilities can be discussed. Additionally, the study director can summarize clinical findings and potential target organs. The prosectors should be familiar with the protocol and amendments for the study involving the animals being necropsied. The protocol should clearly state which tissues are to be collected and weighed and how the tissue should be preserved. During the necropsy, devices such as checklists or pre-labeled

compartmentalized trays should be present to ensure that all required organs are taken and weighed. In recent years commercially available computer software has been developed to assist in the collection and weighing of tissues at necropsy. This is an electronic copy of the checklist, but also provides a method for recording observations in a consistent manner. Copies of the last clinical observation should be present at the necropsy so the prosectors are alerted to lesions that may be present and require special attention. Palpation records are particularly important at carcinogenicity study necropsies to ensure that all masses detected at the last examination are confirmed and collected.

The necropsy will involve a check of animal identification and sex, an external examination of the animal, an *in situ* examination of all tissues and organs (prior to dissection), and the collecting and weighing of the required tissues.

Summary

In summary, there are several advantages to the use of the rat in toxicology studies. Because of its widespread use in many fields of biology, there is a large historical database of information about the anatomy and needs of the species. This knowledge, along with information about the species' metabolism and response to toxicants, has shown the rat to be generally a good model for prediction of the human response to toxicants. Rats have a lifespan of 24–30 months, which is convenient for chronic toxicity and carcinogenicity studies where animals need to be exposed for the majority of their lifetime. The short gestation time and large litter size make the rat a good model for reproductive studies. The development of specific pathogen-free rats and improvement in husbandry has eliminated most of the disease outbreaks that may have introduced variability into a study. The lack of an emetic response allows for the testing of higher dosages of compound that may cause vomiting in other species. The small size of the rat is useful in that large numbers can be housed economically. The size is also useful in that smaller amounts of test compound are required to gain maximal exposure.

The relatively small size of the rat is also one of its major disadvantages. The amount of blood that can be taken from the animal is limited, thus limiting the number of parameters that can be investigated or the number of toxicokinetic samples that can be collected from a single animal. This problem can be overcome by adding additional animals in interim sacrifice groups or by collecting toxicokinetic samples from cohorts of animals at different times. In most cases it is recommended that toxicokinetic samples be collected from satellite animals

and not from the main study animals being used for evaluation of toxicity. However, an increased number of animals means increased work in the conduct of the study. The small size and relatively active nature of the rats makes some procedures, such as intravenous dosing or collection of electrocardiograms, difficult. These issues have been overcome with the use of suitable restrainers or, in some cases, anesthesia. The rat has been used successfully in toxicology research for close to a century and will continue to be used for the foreseeable future.

THE MOUSE

Use in toxicological research

As discussed in earlier sections, the choice of a species for toxicity testing is based on consideration of a range of variables. Ideally, if toxicity testing is intended to provide information on the safety of a test article in or by humans, the species chosen for testing should be most similar to the human in the way it handles the test article pharmacodynamically. Substantial differences in absorption, distribution, metabolism, or elimination between test species and the target species, e.g. the human, will reduce the predictive value of the test results. From a practical standpoint, often the pharmacokinetics are unknown in humans or the variety of available test species at the time of species selection. For this reason, testing is usually conducted in at least two species. Generally, one of those species is usually a rodent and one a nonrodent. The two most commonly used rodent species are mice and rats, and often toxicity testing is conducted in both of those species.

Mice have many advantages as test animals for toxicity testing. They are small, relatively economical to obtain, house, and care for, and they are generally easy to handle. Mice are generally more economical than rats in these respects. While mice may attempt to escape or bite handlers, with regular, gentle handling they are easily managed. Other advantages of the species include a short gestation period and a short natural life span. These characteristics allow studies that include evaluation of reproductive performance or exposure to a test article for periods approaching the expected life span (e.g. evaluation of carcinogenic potential) to be conducted in a practical time frame. High quality, healthy mice are available from reliable commercial suppliers. Many genetically well-defined, highly inbred, specifically or randomly outbred strains are available. Mice have been used in biomedical research for hundreds of years, and because of this, many technical procedures have been developed for use with the species, and a

vast body of historical data is available for most strains. This historical database includes information on optimal nutritional and housing requirements in addition to data such as the expected background incidence of various diseases and types of tumors in untreated animals, and is continually being added to (Blackwell *et al.*, 1995).

There are also disadvantages to using mice, and most are related to the small size of the animal and the limits that this imposes. The smaller size and higher metabolic rate compared to the rat renders the species a bit less hearty than rats. Deviations in environmental conditions such as an air-conditioning failure or failure in an automatic watering system typically have more severe effects on the smaller species such as mice than the same deviations have on rats. Owing to their high level of natural activity, most strains of mice will not become as docile or easy to handle as rats that have received equivalent handling. Small size often precludes or renders more difficult a number of procedures that are commonly conducted in toxicity testing, such as the collection of large samples or repeated samples of blood and urine, electrocardiographic evaluation, and some necropsy evaluations. The FDA provides HED (Human

TABLE 2.17 Normal physiological values general and reproductive

General	
Life span	
average	1–3 years
maximum reported	4 years
Adult weight	
male	20–40 g
female	18–40 g
Surface area	0.03–0.06 cm ²
Chromosome number (diploid)	40
Food consumption	4–5 g/day
Water consumption	5–8 mL/day <i>ad libitum</i>
Body temperature	36.5°C
oxygen consumption	1.69 mL/g/hr
Reproductive	
Age, sexual maturity	
male	50 days (20–35 g)
female	18–40 g
Breeding season	continuous, cyclic
Estrus cycle	4–5 days
Gestation period	
average	19 days
range	17–21 days
Litter size	
average	12
range	1–23
Birth weight	1.5 g
Age begin dry food	10 days
Age at weaning	16–21 days (10–12 g)

Source: Data derived from Jacoby and Fox (1984).

Equivalent Dose) interspecies conversion factors for converting most animal model NOAELs or such to a human equivalent. The conversion factor for the mouse is 12.1, meaning an observed NOAEL in a mouse of 12.1 mg/kg/day would be converted to 1.00 mg/kg/day for conservative human administration. This relatively high value imposes a *de facto* penalty on using mice as one of the species in a safety assessment.

This section will provide brief summaries of some of the normal physiological values and salient features of the species and some of the specific strains that may be useful in selecting an appropriate species and strain for toxicity testing.

Normal physiological values

Selected normal physiological values for mice are shown in Tables 2.17 and 2.18. Median survival of a number of groups of Charles River CD-1 outbred mice is shown in Table 2.19.

These normal values will vary depending upon the strain of mouse, supplier, condition at arrival, type of feed, environmental and housing conditions, and in some cases, time of year. These data should be considered as a reference, but will not necessarily represent experience in any particular laboratory.

TABLE 2.18 Normal physiological values cardiovascular and respiratory

Cardiovascular	
Heart rate	
average	600/min
range	320–800/min
Blood pressure	
Systolic	113–160 mm Hg
Diastolic	102–110 mm Hg
Blood volume	
plasma	45 mL/kg
whole	78 mL/kg
Hematocrit	41.5%
RBC life span	20–30 days
RBC diameter	6.6 microns
Plasma pH	7.2–7.4
Respiratory	
Rate	
average	163/min
range	320–800/min
Tidal volume	
average	0.18 mL
range	0.09–0.38 mL
Minute volume	
average	24 mL/min
range	11–36 mL/min

Source: Data derived from Jacoby and Fox (1984), and from the Animal Diet Reference Guide, Purina Mills, Inc. (Ralston Purina Company, 1987).

Species differences

Mice are similar to other common laboratory animal species and to humans in many ways, yet the differences should not be underestimated. Mice have a high metabolic rate compared to other species. This fact alone may result in increased or decreased toxicity of a test article, depending on the specific mechanism of intoxication. In many cases, high metabolic rate may be associated with rapid absorption, distribution, metabolism, and elimination of a test article. Mice are obligate nose breathers, and have more convoluted nasal passages than humans. This may result in an excess of respirable test article deposited in the nasal passages, resulting in either increased or decreased relative toxicity, depending upon the most critical site of absorption. The small size of the mouse compared to other common laboratory species offers a significant advantage if the test article is expensive or in short supply. As an approximation, a mouse weighs about 10% as much as a rat, about 5% as much as a guinea pig, about 1% as much as a rabbit, and less than 1% as much as a dog or primate. Material requirements to administer equivalent dose levels are usually proportional to body weight, so the test article savings associated with the mouse are evident. The small size of a mouse results in high surface area to body mass ratio, which in turn causes the mouse to be relatively intolerant of thermal and water balance stresses. The kidneys of a mouse have about

twice the glomerular filtering surface per gram of body weight as a rat, and owing to the specific architecture of the murine kidney, they are capable of producing urine that is about four times as concentrated as the highest attainable human concentrations (Jacoby and Fox, 1984). These characteristics of renal architecture and function may be important to the toxicity of some test articles. Mice differ from most species by the formation of a persistent vaginal plug after mating. The presence of a vaginal plug is easily detected, is considered evidence of mating, and is a useful characteristic during the conduct of reproductive studies.

It is also frequently the case in pharmaceutical research and development that the nonclinical efficacy model for a new drug is based on the mouse, making it the natural choice for rodent evaluation of the drug.

Strain differences

In addition to differences between mice and other species, there are important differences among different strains of mice. The appropriate choice of a strain of mice for a particular toxicity study should consider the specific objectives of the study, and the specific characteristics of candidate strains that might assist in or hinder achieving those study objectives.

One difference among strains lies in the normal body weights of various strains at different ages. These differences are summarized for selected strains available from the Charles River Breeding Laboratories in Table 2.20. Outbred strains tend to be larger at maturity than inbred strains, with the CD-1 strain reaching the highest mean weights at 56 days of age of those strains shown in Table 2.20. The CF1 strain has been reported to be highly resistant to mouse typhoid and to be relatively resistant to salmonellosis (Hill, 1981). Nude or athymic strains of mice are more sensitive to tumor development than heterozygous strains. These sensitive strains develop the same types of tumors as those seen in more conventional strains, but the incidences are higher and the latency periods shorter. There is a wide spectrum of

TABLE 2.19 Median survival of 16 groups of control mice (%)

Sex	Period of Time on Study (months)			
	6	12	18	21
Male	98%	91%	63%	46%
Female	98%	95%	74%	68%

Source: Adapted from Lang (1989a,b).

Data represent median survival of Charles River CD-1 outbred albino mice enrolled in 24-month chronic toxicity studies at pharmaceutical or contract toxicology laboratories.

TABLE 2.20 Normal body weights in grams of selected strains of mice

Age (days)	Outbred Strains				Inbred Strains						Hybrid			
	CD-1		CF-1		CFW		C3H		C57BU6		BALB/c		B6C3FI	
	M	F	M	F	M	F	M	F	M	F	M	F	M	F
21	12	11	12	11	9	9	—	—	—	—	—	—	—	—
28	20	18	18	17	16	13	17	16	14	13	16	14	16	14
35	27	22	24	21	19	17	18	17	17	14	17	16	20	17
42	30	24	27	22	24	20	20	18	19	16	18	17	22	18
49	33	26	28	24	27	22	24	23	21	17	20	18	24	19
56	35	27	30	26	28	23	27	26	22	18	21	19	26	21

Source: Data derived from Charles River Growth Charts.

susceptibility to spontaneous lung tumors in various strains of mice, and evidence suggests that there is a high correlation between spontaneous incidence and chemical inducibility in those various strains (Shimkin and Stoner, 1975).

The inbred strain A mouse appears to be the most susceptible to lung tumors, and forms the basis of a lung tumor bioassay, with tumors inducible within 8 weeks or less of treatment. Susceptibility of various strains to the initiation and/or promotion of skin tumors has also been shown to differ greatly (Chouroulinkov *et al.*, 1988; Steinel and Baker, 1988). The incidences of selected spontaneously occurring neoplastic lesions in CD-1 (outbred) and B60171 (hybrid) strains are compared in Table 2.21.

The number of strain-related differences in susceptibility to various test articles and environmental conditions exceeds the scope of this chapter, but additional information is available.

Study designs

Most toxicity and teratology studies conducted in mice are designed to provide information on potential human toxicity. Test substances are typically administered by the expected route of human exposure. A pharmaceutical product that is intended for oral administration (tablet, capsule, solution, or suspension), or a food additive would generally be administered by the oral route. Oral administration to mice is usually accomplished by administration of a solution or suspension by oral gavage, by mixture of the test substance with the diet, or less commonly added to the drinking water.

The specific design of toxicity studies should be tailored to the objective to be achieved and to any specific characteristics of the test substance. Many features of the study design will be predicated on guidelines and practices of regulatory agencies such as the Food and Drug Administration (FDA) or Environmental Protection Agency (EPA) in the United States or their counterparts in other countries to which the results of the 4902MIN are submitted in support of a safety claim. Recommendations for study length (duration of dosing) fall in this category.

Toxicity studies are usually conducted in order of increasing duration of dosing, beginning with acute toxicity studies. When this regimen is followed, each study provides progressively more useful information for the selection of doses for the next, longer study.

Acute toxicity studies

Acute toxicity studies are conducted to evaluate the effects of a single substance. Usually each animal

TABLE 2.21 Incidence of spontaneously occurring neoplastic lesions. Lesions Occurring at spontaneous incidence of ~:1% in either sex of charles river CD-1 or 136C3171 Mice Strain

Location & Lesion	CD-1		B6C3F1	
	M	F	M	F
Lymphoreticular tumors				
lymphosarcoma			6.0	12.0
lymphocytic leukemia			1.3	1.4
lymphoma	3.7	9.9		
histiocytic lymphoma			0.5	1.4
histiocytic sarcoma			0.1	1.4
lymphoblastic lymphoma	1.1	1.7	0.2	0.1
lymphocytic lymphoma	2.6	1.7	0.6	1.7
reticulum cell sarcoma	2.6	5.1	0.4	0.2
Skin/subcutis				
fibrosarcoma	0.2	0.5	1.0	0.5
Mammary gland				
adenocarcinoma		1.7		0.9
Lung				
bronchiolar/alveolar adenoma	4.0	2.9	8.3	3.3
bronchiolar/alveolar carcinoma	3.5	3.1	1.9	0.6
alveolar type II carcinoma	11.7	13.9	0.2	0.1
alveolar type II adenoma			2.5	1.2
adenoma			1.2	0.7
Liver				
nodular hepatocellular prolif.	5.4	1.7	0.5	0.1
hepatocellular adenoma	5.6	0.8	17.2	7.1
hepatocellular carcinoma	7.3	1.0	13.2	2.4
hemangioma	1.0	1.2	0.7	0.3
hemangiosarcoma	1.0	0.2	0.5	0.1
Reproductive system				
Ovary				
cystadenoma		1.1		0.3
Uterus				
endometrial stromal polyp		3.3		2.9
endometrial sarcoma		1.9		0.6
leiomyoma		1.0		
leiomyosarcoma		1.0		0.3
hemangioma		1.0		0.9
Pituitary				
adenoma		3.4	0.3	7.9
Thyroid gland				
follicular cell adenoma	0.2		0.8	2.4
Adrenal				
cortical adenoma	8.6	1.0	0.4	0.3
Harderian gland				
cystadenoma			1.9	1.6
adenoma			1.5	0.6

Source: Data from Charles River Breeding Laboratories, compiled from control animals on 24-month studies completed between 1978 and 1986.

receives a single dose of the test substance in this study design. On rare occasion, repeated doses may be administered, but in any event, all doses are administered within 24 hours or less. Historically, a primary objective of acute toxicity testing was to determine an LD50 dose, or that dose which would be lethal to 50% of the animals treated. To achieve this objective, groups of mice,

often numbering 10 or more per sex, are treated with a single dose of the test substance. Depending upon the rate of survival in the initial group(s), additional groups are added to the study at higher and/or lower doses such that most animals die that receive the highest doses and most survive that receive the lowest doses. Survival is assessed at some predetermined interval after dosing, usually 7 or 14 days, but occasionally as early as 24 hr. The resultant dose-response data can be analyzed by a statistical method such as probit analysis (Finney, 1971) to provide an estimate of the median lethal dose (LD50) and some measure of the precision of that estimate, such as the 95% fiducial limits. There are very few scientifically valid reasons to include determination of the LD50 as a significant objective of acute toxicity testing. Most regulatory agencies have dropped their requirements for a specific value for the LD50, and animal welfare considerations preclude the use of the large numbers of animals previously required.

A more contemporary design for acute toxicity testing attempts to derive a maximum amount of information from a minimum number of animals. Study objectives include determination of the most important clinical signs attributable to high doses of the test substance, time of onset and remission of those signs, possible determination of a minimum lethal dosage, and in the event of lethality, the sequence and timing of effects leading to death or recovery. These objectives are achieved by means of a comprehensive schedule of animal observations following dosing. These objectives can usually be achieved by treating from one to three groups of three to five mice/sex/group at different doses.

Traditionally, acute toxicity testing of potential new pharmaceutical products is conducted in at least three species, with one being a nonrodent, and by at least two routes of administration, one of which is the intended clinical route. Mice are the most frequently selected rodent species for acute toxicity testing. The choice of routes of administration depends on the intended clinical route and on how much is already known about the oral bioavailability of the test substance. If the intended clinical route is oral, acute testing by oral gavage with a solution or suspension is of primary importance. If other clinical routes are anticipated (e.g. intravenous [IV] or dermal), they represent good secondary routes for acute testing. Ordinarily, at least one parenteral route is used for acute testing, and that route may be IV if the product is soluble in a fairly innocuous vehicle (e.g. water or saline) or intraperitoneal (IP) as a suspension if the product is insoluble in an aqueous (or other innocuous) vehicle. If the intended clinical route is not oral, the oral route is usually selected as a secondary route for acute toxicity testing to provide information

relevant to accidental oral ingestion. A rough estimate of oral bioavailability can be based on a comparison of the acute toxicity associated with various doses administered by the oral and parenteral routes. Acute toxicity testing conducted for other purposes is usually more limited in scope. Most regulatory agencies no longer require a full complement of species and routes of administration to render decisions on acute toxicity.

There are a few characteristics of acute toxicity testing that are not common in other toxicity protocols. In a typical repeated dose toxicity study, several groups of animals are treated concurrently with predetermined doses of test substance and a control substance. To reduce animal use in acute toxicity testing, studies that include more than one dose group are usually dosed sequentially, with an interval of at least 24 hrs between dosing of subsequent groups. This allows the effects of the previous dose to be fully manifested, and allows selection of the subsequent dose to provide the highest probability of contributing more useful information. Another unusual aspect of acute toxicity studies is the nutritional status of the animals at dosing. Because some schools believe that the results of acute toxicity testing are more reliable if all animals are in a uniform nutritional state, mice to be dosed orally are often fasted overnight prior to dosing. Fasting allows dose volumes to be higher than in repeated dose studies, and because dosing only occurs on 1 day, dietary stress is considered tolerable. The scientific merits of this practice are debatable, but fasting is "traditional" in oral acute toxicity studies. While the practice of conducting gross necropsies at the end of acute toxicity studies is growing in popularity, this practice rarely yields useful information. The toxicity resulting from acute exposure is usually associated with a biochemical or functional imbalance rather than with a change in the gross or microscopic architecture of an organ system. Changes observable at gross necropsy are more often associated with repeated dosing at sublethal levels. For similar reasons, microscopic examination of tissue is rarely conducted in acute toxicity studies unless there is some scientific reason to expect it would be useful.

The results of a well-designed acute toxicity study can help to predict likely target organ systems and possible outcome in the event of massive human overexposure, can help in establishing risk categories for EPA or Department of Transportation (DOT) classification, and can help in dose selection for the initial repeated dose toxicity tests to be conducted. An example of an acute toxicity study design is in [Table 2.22](#).

Short-term toxicity studies

The objective of short-term or subchronic toxicity studies is to describe and define the toxicity associated with